

Another Boozer-coordinate-free motivation for quasisymmetry

We would like any confinement device to have the following omnigenity property: the ψ coordinate of each trapped particle has no net change per bounce. Equivalently, $\Delta\psi = 0$ where

$$\Delta\psi = \psi_{final} - \psi_{initial} = \oint \frac{d\psi}{dt} dt = \oint \mathbf{v}_d \cdot \nabla \psi dt. \quad (1)$$

Here, \oint indicates an integral over a bounce, holding the magnetic moment $\mu = mv_\perp^2 / (2B)$ and energy $mv^2 / 2$ fixed. Rather than integrate over the actual orbit (which has a nonzero width), we take (1) to be an integral along a field line, which is the same aside from a correction of order ρ_s , and which is much easier to work with mathematically.


Before considering stellarators, let us examine in detail why (1) is satisfied in a tokamak. First, we notice

$$\Delta\psi = \oint (\mathbf{v}_d \cdot \nabla \psi) dt = 2 \int_{\ell_-}^{\ell_+} (\mathbf{v}_d \cdot \nabla \psi) \frac{d\ell}{|v_\parallel|} \quad (2)$$

where ℓ_- and ℓ_+ are the two bounce points. Changing the variable of integration to $B = |\mathbf{B}|$,

$$\Delta\psi = 2 \sum_\gamma \int_{B_{\min}}^{B_{tr}} (\mathbf{v}_d \cdot \nabla \psi) \frac{dB}{|v_\parallel| |\mathbf{b} \cdot \nabla B|} = 2 \sum_\gamma \gamma \int_{B_{\min}}^{B_{tr}} (\mathbf{v}_d \cdot \nabla \psi) \frac{dB}{|v_\parallel| |\mathbf{b} \cdot \nabla B|} \quad (3)$$

where $B_{tr} = mv^2 / (2\mu)$ is the value of B at which the particle is trapped, and where $\gamma = \text{sign}(\mathbf{b} \cdot \nabla B)$ is the “branch”: $\gamma = +1$ if our location relative to B_{\min} is *parallel* to \mathbf{B} , and $\gamma = -1$ if our location relative to B_{\min} is *antiparallel* to \mathbf{B} . The sum over γ is needed because each value of B could correspond to one of two different ℓ , as shown by the following figure. As with any change of variables, the *absolute value* of the Jacobian arises, and in the case of (3) this is $1/|\mathbf{b} \cdot \nabla B|$.



Next, using $\mathbf{B} \times \boldsymbol{\kappa} \cdot \nabla \psi = B^{-1} \mathbf{B} \times \nabla B \cdot \nabla \psi$ (which is true for *any* MHD equilibrium), we obtain

$$\Delta\psi = \frac{2mc}{Ze} \sum_{\gamma} \gamma \int_{B_{\min}}^{B_{\text{tr}}} \left(v_{\parallel}^2 + \frac{v_{\perp}^2}{2} \right) \frac{1}{|v_{\parallel}| B^2} \frac{\mathbf{B} \times \nabla B \cdot \nabla \psi}{\mathbf{B} \cdot \nabla B} dB. \quad (4)$$

In an axisymmetric MHD equilibrium, $\mathbf{B} = \nabla\phi \times \nabla\psi + F(\psi)\nabla\phi$ where ϕ is the toroidal angle and $F(\psi) = RB_{\text{tor}}$ is a flux function. Therefore $\mathbf{B} \times \nabla\psi = F\mathbf{B} - R^2 B^2 \nabla\phi$, so

$$\frac{\mathbf{B} \times \nabla B \cdot \nabla \psi}{\mathbf{B} \cdot \nabla B} = -F(\psi). \quad (5)$$

Then we can pull this factor outside the integral in (4):

$$\Delta\psi = -\frac{2mcF}{Ze} \sum_{\gamma} \gamma \int_{B_{\min}}^{B_{\text{tr}}} \left(v_{\parallel}^2 + \frac{v_{\perp}^2}{2} \right) \frac{1}{|v_{\parallel}| B^2} dB. \quad (6)$$

It turns out that the remaining integral can actually be done, using the relation $v^2 = v_{\parallel}^2 + (2\mu B/m)$ and recalling that μ and ν are fixed. However, we do not even need to do the integral, because the result evidently cannot depend on the branch (only on B_{\min} , B_{tr} , ν , and μ), and so the result will vanish in the γ sum. Physically, the radial drift in the $\gamma > 0$ branch is equal and opposite to the radial drift in the $\gamma < 0$ branch.

In the above argument, we only used axisymmetry in one small way: writing the left-hand side of (5) as a flux function, so that function could be pulled outside the integral. It did not even matter that the ratio equaled RB_{tor} . In fact, it did not even matter that the ratio was a flux function – only that it was branch-independent! Thus, if we can find any *nonaxisymmetric* field in which

$$\frac{\mathbf{B} \times \nabla B \cdot \nabla \psi}{\mathbf{B} \cdot \nabla B} = Y(\psi, B) \quad (7)$$

for some function Y , then (4) would become

$$\Delta\psi = \frac{2mc}{Ze} \sum_{\gamma} \gamma L(B_{\min}, \nu, \mu) \quad (8)$$

where

$$L(B_{\min}, \nu, \mu) = \int_{B_{\min}}^{B_{\text{tr}}} \left(v_{\parallel}^2 + \frac{v_{\perp}^2}{2} \right) \frac{1}{|v_{\parallel}| B^2} Y(\psi, B) dB. \quad (9)$$

Then the sum over γ in (8) immediately vanishes because L is independent of γ . Thus, any magnetic field with the property (7) is omnigenous: $\Delta\psi = 0$.

As shown in Ref. [1], the quasisymmetry condition $B = B(\psi, M\theta - N\zeta)$ for any integers M and N is satisfied if and only if

$$\frac{\mathbf{B} \times \nabla B \cdot \nabla \psi}{\mathbf{B} \cdot \nabla B} = Y(\psi). \quad (10)$$

This condition, that the ratio is a flux function, is a stricter condition than (7). Therefore quasisymmetry implies omnigenity.

I'm not sure if (7) is as general as omnigenity: here we have only proved that (7) is sufficient for omnigenity, not that (7) is necessary.

[1] Helander and Simakov, **PRL** 101, 145003 (2008).