

**Some equations and formulae that might or might not be useful:**

$$\Delta E = h\nu; \nu = \frac{c}{\lambda}; \lambda = \frac{h}{p}$$

$$h = 6.62 \times 10^{-34} \text{ J s}; \hbar = h/(2\pi)$$

$$c = 3 \times 10^8 \text{ m/s}; k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$\hat{p}_x = -i\hbar \frac{\partial}{\partial x}; E_{kinet} = \frac{p^2}{2m}$$

$$\Delta p_x \cdot \Delta x \geq \frac{\hbar}{2};$$

$$E = \frac{\hbar^2 k^2}{2m}; \Psi_+(x) = Ae^{ikx}; \Psi_-(x) = Ae^{-ikx}$$

$$m_e = 9.1 \times 10^{-31} \text{ kg}$$

$$E_n = \frac{h^2 n^2}{8ma^2}; \Psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{\pi n}{a} x\right)$$

$$m_p = 1.67 \times 10^{-27} \text{ kg}$$

$$E_n = \left(n + \frac{1}{2}\right)h\nu; \Psi_n(x) = A_n H_n(\sqrt{\alpha} x) e^{-\alpha x^2/2};$$

$$\nu = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}}; \alpha = \sqrt{k\mu} / \hbar;$$

$$H_0(\xi) = 1; H_1(\xi) = 2\xi; H_2(\xi) = 4\xi^2 - 2;$$

$$H_3(\xi) = 8\xi^3 - 12\xi; \text{ here } \xi = \sqrt{\alpha} x$$

$$A_n = \frac{1}{\sqrt{2^n n!}} \left(\frac{\alpha}{\pi}\right)^{1/4};$$

$$E_{m_l} = \frac{\hbar^2}{2I} m_l^2; \Psi_{m_l}(\phi) = \frac{1}{\sqrt{2\pi}} e^{im_l \phi}; I = \mu r^2;$$

$$\hat{l}_z = -i\hbar \frac{\partial}{\partial \phi}; l_z = m_l \hbar;$$

$$e^{i\alpha} = \cos \alpha + i \sin \alpha;$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha;$$

$$\cos \alpha = (e^{i\alpha} + e^{-i\alpha})/2; \sin \alpha = (e^{i\alpha} - e^{-i\alpha})/2i$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha;$$

$$2 \sin^2 \alpha = 1 - \cos 2\alpha; 2 \cos^2 \alpha = 1 + \cos 2\alpha$$

$$\int_0^\infty e^{-ax^2} x^{2n} dx = \sqrt{\frac{\pi}{a}} \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2^{n+1} a^n}; \text{ (for } n \geq 0)$$

$$\int_0^\infty e^{-ax^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}}; \int_0^\infty e^{-ax^2} x^2 dx = \frac{1}{4a} \sqrt{\frac{\pi}{a}}$$

$$\int_0^a \sin^2\left(\frac{\pi n}{a} x\right) dx = \frac{a}{2}; \int_0^\infty x^n e^{-ax} dx = \frac{n!}{a^{n+1}};$$

I might add more equations