

# Hermitian forms for $Sp(4, \mathbb{R})$

Jeffrey Adams and Annegret Paul

April 8, 2012

## 1 Notation for $Sp(4, \mathbb{R})$

Basic references are Vogan's notes on  $Sp(4, \mathbb{R})$ : [?] (branching) and [?] (Hermitian forms). In fact these notes are largely a rewriting of [?] in more explicit atlas terms.

$G = Sp(4, \mathbb{R})$  with all the usual notation, including **atlas** stuff.

Write  $(x, y)$  for the usual coordinates, in which  $\rho = (2, 1)$ . The *software* coordinates (assuming we define  $G$  as **sc**) are fundamental weight coordinates. Write  $[a, b]$  for these coordinates. The changes of coordinates are

$$(1.1) \quad \begin{aligned} (x, y) &\rightarrow [x - y, y] \\ (a + b, b) &\leftarrow [a, b] \end{aligned}$$

For example  $\rho = (2, 1) = [1, 1]$ .

## 2 Standard Modules

We always write  $H$  for the once-and-for-all fixed Cartan of  $G$ , and  $X^* = X^*(H)$ ,  $X_* = X_*(H)$ . We may also have use for  $H^\vee$ ,  $X^{\vee,*} = X^*(H^\vee)$  and  $X_*^\vee = X_*(H^\vee)$ . We also fixed once and for all a set of positive roots.

In **atlas** language a *parameter* is a triple  $(x, \lambda, \nu)$  where:

- (1)  $x$  is a **kgb** element, set  $\theta = \theta_x$ ;
- (2)  $\lambda \in (X^* + \rho)/(1 - \theta)X^*$ ;
- (3)  $\nu \in (X^* \otimes_{\mathbb{Z}} \mathbb{Q})^{-\theta} = \mathfrak{a}(\mathbb{Q})^*$ .

This defines a standard module  $I(x, \lambda, \nu)$ . The infinitesimal character is

$$(2.1) \quad \gamma = \frac{1}{2}(1 + \theta)\lambda + \frac{1}{2}(1 - \theta)\nu \in X^* \otimes \mathbb{Q}.$$

For example  $\lambda = \nu$  gives  $\gamma = \lambda$  (note that  $\gamma$  is integral).

Recall  $\theta$  defines positive imaginary and real roots, and  $\rho_r, \rho_i$  and  $\rho_{cx}$ .

Some conditions are:

- (a) The parameter is *standard* if  $\langle \lambda, \alpha^\vee \rangle \geq 0$  for all positive imaginary roots.
- (b) The parameter is *final* if for all positive real roots,  $\langle \nu, \alpha^\vee \rangle = 0$  implies  $\langle \lambda + \rho_r, \alpha^\vee \rangle$  is even. Equivalently:  $\langle \lambda, \alpha^\vee \rangle$  is *odd* for all real-simple roots (simple roots in the subsystem of real roots).
- (c) We say  $(x, \lambda, \nu) \equiv (x, w(\lambda + \rho_r) - \rho_r, w\nu)$  for  $w \in W_r$ . Using this we can assume  $\langle \nu, \alpha \rangle \geq 0$  for all positive real roots.
- (d) If  $\alpha$  is simple and  $\theta_x$ -complex we say  $(x, \lambda, \nu) \equiv (s_\alpha \times x, s_\alpha \lambda, s_\alpha \nu)$ . This allows us to move  $x$  to any fiber on the same Cartan.
- (e) For a standard parameter,  $I(x, \lambda, \nu)$  is zero if and only if  $\langle \lambda, \alpha^\vee \rangle = 0$  for some imaginary-simple root which is compact.

A nonzero final standard module  $I(x, \lambda, \nu)$  has a unique irreducible quotient  $J(x, \lambda, \nu)$ .

We generally write the subscript  $K$  to indicate restriction to  $K$ ; so here we have  $I_K(x, \lambda, \nu)$  and  $J_K(x, \lambda, \nu)$ .

Recall each fiber has a distinguished basepoint, and each conjugacy class of fibers has a canonical fiber. In the output of KGB, the canonical fiber is labelled #, and the basepoints have entry  $(0, \dots, 0)$  preceding the #.

Using (c) we can write every final standard parameter in the form  $(x, \lambda, \nu)$  where  $x$  is in a canonical fiber.

## 2.1 Standard Modules for $Sp(4, \mathbb{R})$

There are 11 kgb elements numbered 0 – 10. I'll call them  $x_0, \dots, x_{10}$ . Here is the output of KGB.

|    |   |       |   |   |   |   |         |     |
|----|---|-------|---|---|---|---|---------|-----|
| 0: | 0 | [n,n] | 1 | 2 | 4 | 5 | (0,0)#0 | e   |
| 1: | 0 | [n,n] | 0 | 3 | 4 | 6 | (1,0)#0 | e   |
| 2: | 0 | [c,n] | 2 | 0 | * | 5 | (0,1)#0 | e   |
| 3: | 0 | [c,n] | 3 | 1 | * | 6 | (1,1)#0 | e   |
| 4: | 1 | [r,C] | 4 | 9 | * | * | (0,0)   | 1 1 |
| 5: | 1 | [C,r] | 7 | 5 | * | * | (0,0)   | 2 2 |

|     |   |       |    |    |    |    |         |         |   |
|-----|---|-------|----|----|----|----|---------|---------|---|
| 6:  | 1 | [C,r] | 8  | 6  | *  | *  | (1,0)   | 2       | 2 |
| 7:  | 2 | [C,n] | 5  | 8  | *  | 10 | (0,0)#2 | 1,2,1   |   |
| 8:  | 2 | [C,n] | 6  | 7  | *  | 10 | (0,1)#2 | 1,2,1   |   |
| 9:  | 2 | [n,C] | 9  | 4  | 10 | *  | (0,0)#1 | 2,1,2   |   |
| 10: | 3 | [r,r] | 10 | 10 | *  | *  | (0,0)#3 | 2,1,2,1 |   |

The basepoints in the canonical fibers are  $x_0, x_7, x_9, x_{10}$ .

**Cartan 0** This is the compact Cartan subgroup, so  $\nu = 0$ , and  $\lambda \in X^* \simeq \mathbb{Z}^2$ . The standard modules are  $(a, b \in \mathbb{Z}, a \geq b \geq 0)$ :

$$\begin{aligned}
I(x_0, (a, b)) & \quad (\text{large DS}) \\
I(x_1, (a, b)) & \quad (\text{large DS}) \\
I(x_2, (a, b)) & \quad (\text{holomorphic DS}) \\
I(x_3, (a, b)) & \quad (\text{antiholomorphic DS})
\end{aligned}$$

These are always final. The standard modules are always nonzero in the first two cases, and if and only if  $a > b$  in the last two.

**Cartan 1** This is the  $\mathbb{C}^*$  Cartan subgroup. The canonical fiber has  $\theta = w = 2, 1, 2$ , so  $\theta(x, y) = (-y, -x)$ . Then  $(1 - \theta)X^* = \{(c, c) \mid c \in \mathbb{Z}\}$  so  $\lambda = (a, b) \bmod (c, c)$  with  $a, b, c \in \mathbb{Z}$ . (This is isomorphic to  $\mathbb{Z}$  by the map  $(a, b) \rightarrow a - b$ ). On the other hand  $(X^* \otimes \mathbb{Q})^{-\theta} = \{(x, x) \mid x \in \mathbb{Q}\}$ .

The root  $(1, -1)$  is imaginary, and  $(1, 1)$  is real. The standard limit modules are

$$(2.1.2)(a) \quad I(x_9, (a, b) \bmod (c, c), (x, x)) \quad (a \geq b, x \geq 0)$$

with infinitesimal character

$$\gamma = \left(x + \frac{a-b}{2}, x - \frac{a-b}{2}\right)$$

These standard modules are always nonzero.

In this case  $M \simeq GL(2, \mathbb{R})$ . The standard module given by  $a, b, x$  is the one associated to the representation on  $M$  with restriction to  $SL(2, \mathbb{R})^\pm$  the discrete series with infinitesimal character  $a - b = 0, 1, 2, \dots$ , and such that  $\text{diag}(e^t, e^t)$  acts by the scalar  $e^{2tx}$ . (These may be not quite the inducing data since we are working with the “ $\Lambda$ -parameters”; see [?] or [?].)

The final condition is

$$x = 0 \Rightarrow a - b \in 2\mathbb{Z} + 1.$$

Suppose  $r \geq s \geq 0$ . Provided  $r, s \in \mathbb{Z}$  there are two standard modules with infinitesimal character  $(r, s)$ :

$$(2.1.2)(b) \quad I(x_9, (r-s, 0), \tfrac{1}{2}(r+s, r+s)) \rightarrow \gamma = (r, s)$$

$$(2.1.2)(c) \quad I(x_9, (r+s, 0), \tfrac{1}{2}(r-s, r-s)) \rightarrow \gamma = (r, -s)$$

In these coordinates, the final condition is:

$$\text{If } x = 0 \text{ then } I(x_9, (c, 0), (0, 0)) \text{ is final} \Leftrightarrow c \in 2\mathbb{Z} + 1.$$

In (b)  $\gamma$  is dominant, whereas in (c) it is not if  $s > 0$ . Using the cross action of the complex root  $(0, 2)$  (note that  $\theta_{x_4}(x, y) = (y, x)$ ):

$$(2.1.3) \quad I(x_9, (r+s, 0), \tfrac{1}{2}(r-s, r-s)) = I(x_4, (r+s, 0), \tfrac{1}{2}(r-s, -s+r))$$

It is convenient to change variables, and renormalize by  $\frac{1}{2}$  as follows. Define

$$(2.1.4)(a) \quad I(x_9, c, x) = I(x_9, (c, 0), \tfrac{1}{2}(x, x)) \quad (c \in \mathbb{Z}_{\geq 0}, x \in \mathbb{R}_{\geq 0}).$$

This has infinitesimal character

$$(2.1.4)(b) \quad \gamma = \tfrac{1}{2}(x+c, x-c).$$

**Cartan 2** This is the  $\mathbb{R}^* \times S^1$  Cartan subgroup. The canonical involution is  $1, 2, 1$ , i.e.  $\theta(a, b) = (-a, b)$ . Therefore  $(1 - \theta)X^* = (2\mathbb{Z}, 0)$ , and  $\lambda = (\bar{a}, b) \in \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$ . On the other hand  $\nu = (x, 0)$  with  $x \in \mathbb{Q}$ . The root  $(2, 0)$  is real, and  $(0, 2)$  is imaginary. So the standard modules are

$$\begin{aligned} I(x_7, (\bar{a}, b), (x, 0)) \\ I(x_8, (\bar{a}, b), (x, 0)) \end{aligned}$$

with  $b \in \mathbb{Z}_{\geq 0}, x \in \mathbb{Q}_{\geq 0}$  and  $\bar{a} \in \mathbb{Z}/2\mathbb{Z}$ . The infinitesimal character is

$$\gamma = (x, b)$$

The final condition is

$$x = 0 \Rightarrow \bar{a} = 1.$$

These standard modules are always nonzero.

The standard modules are those given by the holomorphic discrete series of  $SL(2, \mathbb{R})$  with infinitesimal character  $b = 0, 1, 2, \dots$ , and the character of  $\mathbb{R}^* t \rightarrow |t|^x \text{sgn}(x)^a$ . The standard modules  $I(x_8, (\bar{a}, b), (x, 0))$  are the same, with antiholomorphic in place of holomorphic.

If  $x \geq b$  then  $\gamma$  is dominant. If  $x < b$  it makes sense to apply (d) of Section ???. Using the fact that for  $x_5, x_6$ ,  $\theta(x, y) = (x, -y)$ , with the obvious notation we have

$$\begin{aligned} I(x_7, (\bar{a}, b), (x, 0)) &= I(x_5, (b, \bar{a}), (0, x)) \\ I(x_8, (\bar{a}, b), (x, 0)) &= I(x_6, (b, \bar{a}), (0, x)) \end{aligned}$$

**Cartan 3** This is the split Cartan subgroup, with  $x = x_{10}$ ,  $\theta = -1$ ,  $(1 - \theta)X^* = 2\mathbb{Z} \times 2\mathbb{Z}$ ,  $\lambda = (\bar{a}, \bar{b}) \in \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ,  $\nu = (x, y) \in \mathbb{Q}^2$ . The standard modules are

$$I(x_{10}, (\bar{a}, \bar{b}), (x, y)) \quad (x \geq y \geq 0)$$

with infinitesimal character  $(x, y)$ . The *final* condition is

$$\begin{aligned} y = 0 &\Rightarrow \bar{b} = 1 \\ x - y = 0 &\Rightarrow \bar{a} - \bar{b} = 1 \end{aligned}$$

These standard modules are always nonzero.

### 3 nblock

The `nbblock` command is essential to these computations. The translation between `nbblock` coordinates and human ones is tricky. The main point of this section is to explain the `nbblock` command, and give a translation for  $Sp(4, \mathbb{R})$  between `nbblock` and human coordinates.

Here is an overview of the `nbblock` command, which (among other things) allows the user to input a general standard module. Recall a standard module is determined by a parameter  $(x, \lambda, \nu)$ . See Section ??.

The user first inputs a Cartan, from the `atlas` list `0..n`. The software then gives a list of `kgb` elements `x` in the distinguished fiber for this Cartan. In particular this specifies a specific Cartan involution  $\theta$  of  $H$ .

The user chooses one of the elements  $x$ .

Next the user defines  $\lambda$ . The user inputs  $(\lambda - \rho)_{in} \in X^*$  in the “software” coordinates. Suppose  $G$  is semisimple. If it is entered as `sc` these are fundamental weight coordinates, while `ad` gives simple root coordinates. The

software computes

$$\lambda = (\lambda - \rho)_{in} + \rho.$$

Only the image of  $\lambda$  in  $\rho + X^*/(1 - \theta)X^*$  matters.

Next the user defines  $\nu \in (X^* \otimes_{\mathbb{Z}} \mathbb{Q})^{-\theta} = \mathfrak{a}(\mathbb{Q})^*$ . The user inputs an arbitrary element  $\nu_{in} \in X^* \otimes_{\mathbb{Z}} \mathbb{Q}$  (in software coordinates), and the software defines

$$\nu = \frac{1}{2}(1 - \theta)\nu_{in} \in \mathfrak{a}(\mathbb{Q})^*$$

Let

$$\lambda_0 = \frac{1}{2}(1 + \theta)\lambda.$$

Then (cf. (??)) the infinitesimal character is computed as

$$\gamma = \lambda_0 + \nu = \frac{1}{2}(1 + \theta)\lambda + \frac{1}{2}(1 - \theta)\nu_{in}.$$

**NB:** Only  $\nu = (1 - \theta)\nu_{in}$  matters, and for the infinitesimal character only  $\lambda_0 = (1 + \theta)\lambda$  matters. However,  $\lambda$  has extra (torsion) information not contained in  $\lambda_0$ . For example, if the Cartan is split  $\lambda_0 = 0$  but  $\lambda$  is an element of  $\rho + X^*/2X^*$ .

If  $\gamma$  is dominant the triple  $(x, \lambda, \nu)$  defines a standard module. If the infinitesimal character is singular it may be reducible or 0. If  $\gamma$  is not dominant on the integral roots, the software uses complex cross actions to replace this triple with  $(x', \lambda', \nu')$  such that  $\gamma' = \lambda'_0 + \nu'$  is dominant, and conjugate to  $\gamma$ .

## 4 Standard Modules for $Sp(4, \mathbb{R})$ using `nblock`

We show how to construct the standard modules of Section ?? using `nblock`. This is primarily just a change of coordinates.

In each case the user will enter a Cartan, `kgb` element,  $(\lambda - \rho)_{in} = [\tilde{a}, \tilde{b}]$ , and  $\nu_{in} = [\tilde{x}, \tilde{y}]$ .

### (a) Cartan #0

This case is clear. If  $0 \leq k \leq 3$ :

$$I(x_k, (a, b)) = I(x_k, [\tilde{a}, \tilde{b}], [\tilde{x}, \tilde{y}])$$

provided  $a = \tilde{a} + \tilde{b} + 2, b = \tilde{b} + 1$  ( $\tilde{x}, \tilde{y}$  are irrelevant).

### (b) Cartan #1

This is the  $\mathbb{C}^*$  Cartan, with  $M \simeq GL(2, \mathbb{R})$ .

As in Section ??, there is only one kgb element  $\mathbf{x}_9$  in the distinguished fiber,  $\theta = 2, 1, 2$ ,  $\theta(x, y) = (-y, -x)$ . In fundamental weight coordinates  $\theta[x, y] = [x, -x - y]$ .

The coroot  $[1, 0] = (1, -1)$  is imaginary, so the software requires  $\tilde{a} \geq -1$ . Since  $(1 - \theta)X^* = [0, \mathbb{Z}]$ ,  $\tilde{b}$  has no effect (the Cartan is connected).

If  $\nu = [\tilde{x}, \tilde{y}]$  then  $\frac{1}{2}(1 - \theta)\nu = [0, \tilde{y} + \frac{\tilde{x}}{2}]$ . It is convenient to take  $\tilde{x} = 0$ , so  $\frac{1}{2}(1 - \theta)\nu = \nu$ .

So the module

$$I(x_9, (a, b) \bmod(c, c), (x, x)) \quad (a - b \geq 0, x \geq 0)$$

of Section ?? is given in these coordinates by

$$I(x_9, [\tilde{a}, 0] \bmod(0, \mathbb{Z}), [0, \tilde{x}]) \quad (\tilde{a} \geq -1, \tilde{x} \geq 0)$$

with  $\tilde{a} = a - b - 1 \geq -1$  and  $\tilde{x} = x$ . In particular the infinitesimal characters match up:

$$(x + \frac{a - b}{2}, x - \frac{a - b}{2}) = (\tilde{x} + \frac{\tilde{a} + 1}{2}, \tilde{x} - \frac{\tilde{a} + 1}{2}).$$

In terms of the `nblock` interaction, here is what happens (dropping the tildes). Choose  $x_9$ ,

$$\begin{aligned} (\lambda - \rho)_{in} &= [a, b] = (a + b, b) \\ \nu &= \nu_{in} = [0, x] = (x, x) \end{aligned}$$

If  $x \geq \frac{a+1}{2}$  then the software computes:

$$\begin{aligned} \lambda &= [a + 1, b + 1] = (a + b + 2, b + 1) \\ \lambda_0 &= [a + 1, -\frac{a + 1}{2}] = (\frac{a + 1}{2}, -\frac{a + 1}{2}) \\ \gamma &= [a + 1, x - \frac{a + 1}{2}] = (x + \frac{a + 1}{2}, x - \frac{a + 1}{2}) \end{aligned}$$

and  $\gamma$  is dominant.

If  $x < \frac{a+1}{2}$ ,  $\gamma$  is not dominant, so (provided that  $x - \frac{a+1}{2}$  is an integer) the software conjugates  $\lambda, \nu$  by  $s_2$ . The new Cartan involution is  $\theta(x, y) = (y, x)$ ,

and we get:

$$\begin{aligned}\lambda &= [a + 2b + 3, -b - 1] = (a + b + 2, -b - 1) \\ \lambda_0 &= [0, \frac{a+1}{2}] = (\frac{a+1}{2}, \frac{a+1}{2}) \\ \nu &= [2x, -x] = (x, -x) \\ \gamma &= [2x, -x + \frac{a+1}{2}] = (x + \frac{a+1}{2}, -x + \frac{a+1}{2})\end{aligned}$$

**(c) Cartan #2**

This is the Cartan  $S^1 \times \mathbb{R}^*$ , with  $M = SL(2, \mathbb{R}) \times \mathbb{R}^*$ . As in Section ?? the distinguished fiber has  $\theta = 1, 2, 1$ , so  $\theta(x, y) = (-x, y)$ . In fundamental weight coordinates  $\theta[a, b] = [-a - 2b, b]$ . The coroot  $[0, 1] = (0, 1)$  is imaginary, so we have the condition  $\tilde{b} \geq -1$ . Since  $(1 - \theta)X^* = [2\mathbb{Z}, 0]$ , only the image of  $a \in Z/2\mathbb{Z}$  matters, so we could take  $\tilde{a} = 0, 1$ .

Since  $\nu_{in} = [\tilde{x}, \tilde{y}]$  gives  $\nu = \frac{1}{2}(1 - \theta)\nu_{in} = [\tilde{x} + \tilde{y}, 0]$ , we may as well take

$$\nu_{in} = [\tilde{x}, 0] = (\tilde{x}, 0)$$

in which case  $\nu = \nu_{in}$ .

The translation is, with  $k = 7, 8$ ,

$$I(x_k, (\bar{a}, b), (x, 0)) \quad (b, x \geq 0)$$

is given in the new coordinates by

$$I(x_k, [\tilde{a}, \tilde{b}], [\tilde{x}, 0]) \quad (\tilde{b} \geq -1, \tilde{x} \geq 0)$$

provided  $\tilde{a} + \tilde{b} \equiv \bar{a} \pmod{2}$ ,  $\tilde{b} + 1 = b$  and  $\tilde{x} = x$ .

In terms of the software interaction (dropping the tildes), if  $x \geq b + 1$ :

$$\begin{aligned}\nu &= [x, 0] = (x, 0) \\ \lambda &= [a + 1, b + 1] = (a + b + 2, b + 1) \\ \lambda_0 &= [-b - 1, b + 1] = (0, b + 1) \\ \gamma &= [x - b - 1, b + 1] = (x, b + 1)\end{aligned}$$

and  $\gamma$  is dominant.

If  $x < b + 1$   $\gamma$  is not dominant, so we conjugate everything by  $s_1$ . The new Cartan involution is  $s_2 : (x, y) = (x, -y)$ .



$$\begin{aligned}
\lambda &= [-a-1, a+b+2] = (b+1, a+b+2) \\
\lambda_0 &= [b+1, 0] = (b+1, 0) \\
\nu &= [-x, x] = (0, x) \\
\gamma &= [b+1-x, x] = (b+1, x)
\end{aligned}$$

(d) **Split Cartan**

$$I(x_{10}, (\bar{a}, \bar{b}), (x, y))$$

is the same as

$$I(x_{10}, [\tilde{a}, \tilde{b}], [\tilde{x}, \tilde{y}])$$

provided  $\bar{a} = \tilde{a} + \tilde{b} \pmod{2}$ ,  $\bar{b} = \tilde{b} + 1 \pmod{2}$ , and  $\tilde{x} + \tilde{y} = x, \tilde{y} = y$ .

## 5 Standard K-Representations

Suppose  $I(x, \lambda, 0)$  is a nonzero, final standard limit representation, with  $\nu = 0$ . As in Section ?? its restriction to  $K$  is  $I_K(x, \lambda, 0)$ . This is an important object so we give it a name:

**Definition 5.1**  $I_K(x, \lambda)$  is the restriction of the nonzero, final standard limit module  $I(x, \lambda, 0)$  to  $K$ .

These are a basis of the Grothendieck group of  $K$ . Here is a list of these modules for  $Sp(4, \mathbb{R})$ . The last column gives the highest weight of the lowest  $K$ -types.

$$\begin{aligned}
(5.2) \quad & \begin{array}{lll} I_K(x_0, (a, b)) & a \geq b \geq 0 & (a+1, -b) \\ I_K(x_1, (a, b)) & a \geq b \geq 0 & (b, -a-1) \\ I_K(x_2, (a, b)) & a > b \geq 0 & (a+1, b+2) \\ I_K(x_3, (a, b)) & a > b \geq 0 & (-b-2, -a-1) \\ I_K(x_9, (c, 0)) & c > 0 \text{ odd} & (\frac{c+1}{2}, -\frac{c+1}{2}) \\ I_K(x_7, (\bar{1}, b)) & b \geq 0 & (b+1, 1) \\ I_K(x_8, (\bar{1}, b)) & b \geq 0 & (-1, -b-1) \\ I_K(x_{10}, (\bar{0}, \bar{1})) & & (0, 0) \end{array}
\end{aligned}$$

We will need some Hecht-Schmid identities for some non-final parameters. For example  $I(x_9, (c, 0), (0, 0))$  is not final if  $c$  is even. To see this using

the software, use the `Ktypeform` command. In this example  $\lambda = (2, 0)$ ,  $\lambda - \rho = (0, -1)$  which equals  $[1, -1]$  in fundamental weight coordinates.

```
real: Ktypeform
Choose KGB element: 9
2rho = [ 2, 2 ]
Give lambda-rho: 1 -1
Representation [ 4, 4 ]@(0,0)#1 is not final, as witnessed by coroot [1,2].
```

This is because

$$(5.3)(a) \quad I(x_9, c, 0) = J(x_0, \tfrac{1}{2}(c, c)) + J(x_1, \tfrac{1}{2}(c, c))$$

and so

$$(5.3)(b) \quad I_K(x_9, (c, 0)) = I_K(x_0, \tfrac{1}{2}(c, c)) + I_K(x_1, \tfrac{1}{2}(c, c))$$

with lowest  $K$ -types

$$(5.3)(c) \quad (\tfrac{c}{2} + 1, -\tfrac{c}{2}), (\tfrac{c}{2}, -\tfrac{c}{2} - 1)$$

Similarly if  $b \geq 0$ :

$$(5.4)(a) \quad \begin{aligned} I_K(x_7, \bar{0}, b) &= I_K(x_0, (b, 0)) + I_K(x_2, (b, 0)) \\ I_K(x_8, \bar{0}, b) &= I_K(x_1, (b, 0)) + I_K(x_3, (b, 0)) \end{aligned}$$

with lowest  $K$ -types

$$(5.4)(b) \quad \begin{aligned} I_K(x_7, (\bar{0}, b)) & \quad (b+1, 0), (b+1, 2) \\ I_K(x_8, (\bar{0}, b)) & \quad (0, -b-1), (-2, -b-1) \end{aligned}$$

without the second term if  $b = 0$ .

**Question:** What is the best way to see this using the software?

## 6 Reducibility: Real Roots (parity condition)

Suppose  $I(x, \lambda, \nu)$  is a standard module. A real root  $\alpha$  satisfies the *parity* condition if

$$(6.1)(a) \quad \lambda(m_\alpha) = -(-1)^{\langle \nu + \rho_r, \alpha^\vee \rangle}$$

or equivalently

$$(6.1)(b) \quad \langle \lambda + \nu + \rho_r, \alpha^\vee \rangle \in 2\mathbb{Z} + 1.$$

This holds if and only if  $I(x, \lambda, \nu)$  has some reducibility accounted for by  $SL(2, \mathbb{R})_\alpha$ .

**Note** Here is a mnemonic for remembering this condition. The trivial representation of  $SL(2, \mathbb{R})$  is given by the holomorphic character  $\rho$ :  $\rho(z) = z$  ( $z \in \mathbb{C}^*$ ). The character  $\rho$  is given by  $(\lambda, \nu) = (\rho, \rho)$ , and note that  $\rho = \rho_r$ . So (b) says  $\langle 3\rho, \alpha^\vee \rangle \in 2\mathbb{Z} + 1$  which is true; this standard module is reducible.

Note that *no real root satisfies the parity condition* if and only if

$$(6.1)(c) \quad \langle \lambda + \nu, \alpha^\vee \rangle \in 2\mathbb{Z} + 1 \text{ for all real-simple roots } \alpha.$$

In particular

$$(6.1)(d) \quad I(x, \lambda, \nu) \text{ irreducible} \Rightarrow (??)(c) \text{ holds.}$$

An important special case is  $\nu = 0$ :

$$(6.1)(e) \quad I(x, \lambda, 0) \text{ irreducible} \Rightarrow \langle \lambda, \alpha^\vee \rangle \in 2\mathbb{Z} + 1 \text{ for all real-simple roots } \alpha.$$

This is the *final* condition of Section ???. Also see the help file for the `Ktypeform` command.

**Cartan 1** There is a single real root  $\alpha = (1, 1)$ , and the parity condition for  $I(x_9, (a, b) \pmod{(c, c)}, (x, x))$  is:

$$a + b + 2x + 1 \in 2\mathbb{Z} + 1$$

**Cartan 2** The real root is  $(2, 0)$ , and the parity condition for  $I(x_k, (\bar{a}, b), (x, 0))$ ,  $k = 7, 8$ , is

$$\bar{a} + x + 1 \in 2\mathbb{Z} + 1$$

**Cartan 3** All roots are real. The parity conditions on  $I(x_{10}, (\bar{a}, \bar{b}), (x, y))$  are

$$\begin{aligned}
(6.2) \quad & \langle \lambda + \nu + \rho_r, \alpha^\vee \rangle \in 2\mathbb{Z} + 1. \\
& \alpha_1 = (1, -1) : \quad \bar{a} - \bar{b} + x - y + 1 \in 2\mathbb{Z} + 1 \\
& \alpha_2 = (0, 2) : \quad \bar{b} + y + 1 \in 2\mathbb{Z} + 1 \\
& (1, 1) : \quad \bar{a} + \bar{b} + x + y + 3 \in 2\mathbb{Z} + 1 \\
& (2, 0) : \quad \bar{a} + x + 2 \in 2\mathbb{Z} + 1
\end{aligned}$$

For example for  $I(x_{10}, (2, 1), (2, 1))$ ,  $\alpha_1, \alpha_2$  satisfy the parity condition. On the other hand for  $I(x_{10}, (0, 0), (2, 1))$ , no root satisfies the parity condition.

We collect these conditions in a table, using the alternate parametrization of (??) for Cartan 1.

Parity for  $Sp(4, \mathbb{R})$

| Cartan | Module                                                | root    | parity condition                                    |
|--------|-------------------------------------------------------|---------|-----------------------------------------------------|
| 0      | no real roots                                         |         |                                                     |
| 1      | $I(x_9, (r \pm s, 0), \frac{1}{2}(r \mp s, r \mp s))$ | (1, 1)  | $2r + 1 \in 2\mathbb{Z} + 1$                        |
| 2      | $I(x_k, (\bar{a}, b), (x, 0))$                        | (2, 0)  | $\bar{a} + x + 1 \in 2\mathbb{Z} + 1$               |
| 3      | $I(x_{10}, (\bar{a}, \bar{b}), (x, y))$               | (1, -1) | $\bar{a} - \bar{b} + x - y + 1 \in 2\mathbb{Z} + 1$ |
| 3      |                                                       | (0, 2)  | $\bar{b} + y + 1 \in 2\mathbb{Z} + 1$               |
| 3      |                                                       | (2, 0)  | $\bar{a} + x + 2 \in 2\mathbb{Z} + 1$               |
| 3      |                                                       | (1, 1)  | $\bar{a} + \bar{b} + x + y + 3 \in 2\mathbb{Z} + 1$ |

On Cartan 1 recall  $\alpha$  is integral if  $2r \in \mathbb{Z}$ .

It might be better to write the conditions on the split Cartan (the last four entries) as follows:

$$\begin{aligned}
(1, -1) \quad & (\bar{a} + x) - (\bar{b} + y) \in 2\mathbb{Z} \\
(0, 2) \quad & \bar{b} + y \in 2\mathbb{Z} \\
(2, 0) \quad & (\bar{a} + x) \in 2\mathbb{Z} + 1 \\
(1, 1) \quad & (\bar{a} + x) + (\bar{b} + y) \in 2\mathbb{Z}
\end{aligned}$$

## 7 Reducibility: Complex Roots

Suppose  $I(x, \lambda, \nu)$  is standard, with regular infinitesimal character. An integral complex root  $\alpha$  contributes to reducibility of  $I(x, \lambda, \nu)$  if and only if  $\theta_x(\alpha) < 0$ ; equivalently  $\alpha \in \tau(I)$ , or  $\alpha$  is of type  $\mathbf{C^-}$ .

For  $Sp(4, \mathbb{R})$  only Cartans 1, 2 have complex roots.

**Cartan 1** Consider  $I(x_9, (r \pm s, 0), \frac{1}{2}(r \mp s, r \mp s))$  with  $r, s \geq 0, r \pm s \in \mathbb{Z}$ . Assume  $\gamma = (r, \mp s)$  is regular, i.e.  $r > s > 0$ .

Consider the positive complex root  $\alpha = (2, 0)$ . Assume this is integral, i.e.  $r \in \mathbb{Z}$ . Then  $\theta(\alpha) = (0, -2)$ , and this is positive or negative depending on the sign. Thus:

$$\begin{aligned} I(x_9, (r - s, 0), \frac{1}{2}(r + s, r + s)) &\rightarrow \alpha \text{ is type } \mathbb{C}- \\ I(x_9, (r + s, 0), \frac{1}{2}(r - s, r - s)) &\rightarrow \alpha \text{ is type } \mathbb{C}+ \end{aligned}$$

**Cartan 2** Consider  $I(x_k, (\bar{a}, b), (x, 0))$  with  $k = 7, 8$ , and  $x, b \geq 0$ . so  $\gamma = (x, b)$ .

Assume  $\gamma$  is regular, i.e.  $x \neq b$ , and  $x, b \neq 0$ , and  $\alpha = (1, 1)$  is integral, i.e.  $x + b \in \mathbb{Z}$ . Recall  $\theta(a, b) = (-a, b)$ . Thus  $\alpha > 0$  is complex, and  $\theta(\alpha) = (-1, 1) > 0$  if  $x < b$ , and  $< 0$  otherwise. Therefore

$$\alpha \text{ is type } \mathbb{C}- \Leftrightarrow x > b.$$

## 8 Integral blocks of $Sp(4, \mathbb{R})$

Here is the big block:

|           |   |         |    |    |         |         |   |         |
|-----------|---|---------|----|----|---------|---------|---|---------|
| 0( 0,6):  | 0 | [i1,i1] | 1  | 2  | ( 4, *) | ( 5, *) | 0 | e       |
| 1( 1,6):  | 0 | [i1,i1] | 0  | 3  | ( 4, *) | ( 6, *) | 0 | e       |
| 2( 2,6):  | 0 | [ic,i1] | 2  | 0  | ( *, *) | ( 5, *) | 0 | e       |
| 3( 3,6):  | 0 | [ic,i1] | 3  | 1  | ( *, *) | ( 6, *) | 0 | e       |
| 4( 4,5):  | 1 | [r1,C+] | 4  | 9  | ( 0, 1) | ( *, *) | 1 | 1       |
| 5( 5,4):  | 1 | [C+,r1] | 7  | 5  | ( *, *) | ( 0, 2) | 2 | 2       |
| 6( 6,4):  | 1 | [C+,r1] | 8  | 6  | ( *, *) | ( 1, 3) | 2 | 2       |
| 7( 7,3):  | 2 | [C-,i1] | 5  | 8  | ( *, *) | (10, *) | 2 | 1,2,1   |
| 8( 8,3):  | 2 | [C-,i1] | 6  | 7  | ( *, *) | (10, *) | 2 | 1,2,1   |
| 9( 9,2):  | 2 | [i2,C-] | 9  | 4  | (10,11) | ( *, *) | 1 | 2,1,2   |
| 10(10,0): | 3 | [r2,r1] | 11 | 10 | ( 9, *) | ( 7, 8) | 3 | 2,1,2,1 |
| 11(10,1): | 3 | [r2,rn] | 10 | 11 | ( 9, *) | ( *, *) | 3 | 2,1,2,1 |

Fix an integral regular infinitesimal character  $(a, b)$  with  $a > b > 0$ . Here are the parameters realized using only one fiber on each Cartan, using the cross action to replace  $x_4$  with  $x_9$ , and  $x_{4,5}$  with  $x_{7,8}$ :

| number | Cartan | length | x  | $\lambda$                    | $\nu$                                      |
|--------|--------|--------|----|------------------------------|--------------------------------------------|
| 0      | 0      | 0      | 0  | $(a, b)$                     | *                                          |
| 1      | 0      | 0      | 1  | $(a, b)$                     | *                                          |
| 2      | 0      | 0      | 2  | $(a, b)$                     | *                                          |
| 3      | 0      | 0      | 3  | $(a, b)$                     | *                                          |
| 4      | 1      | 1      | 9  | $(a + b, 0)$                 | $(\frac{1}{2}(a - b), \frac{1}{2}(a - b))$ |
| 5      | 2      | 1      | 7  | $(\bar{b}, a)$               | $(b, 0)$                                   |
| 6      | 2      | 1      | 8  | $(\bar{b}, a)$               | $(b, 0)$                                   |
| 7      | 2      | 2      | 7  | $(\bar{a}, b)$               | $(a, 0)$                                   |
| 8      | 2      | 2      | 8  | $(\bar{a}, b)$               | $(a, 0)$                                   |
| 9      | 1      | 2      | 9  | $(a - b, 0)$                 | $(\frac{1}{2}(a + b), \frac{1}{2}(a + b))$ |
| 10     | 3      | 3      | 10 | $(\bar{a}, \bar{b})$         | $(a, b)$                                   |
| 11     | 3      | 3      | 10 | $(\bar{a} + 1, \bar{b} + 1)$ | $(a, b)$                                   |

Note that 10 is the trivial representation, given by  $\lambda = \nu = \rho$ .

Alternatively, using all  $x_i$ , so the  $x$ -values match up with the output of block:

| number | Cartan | length | x  | $\lambda$                    | $\nu$                                       |
|--------|--------|--------|----|------------------------------|---------------------------------------------|
| 0      | 0      | 0      | 0  | $(a, b)$                     | *                                           |
| 1      | 0      | 0      | 1  | $(a, b)$                     | *                                           |
| 2      | 0      | 0      | 2  | $(a, b)$                     | *                                           |
| 3      | 0      | 0      | 3  | $(a, b)$                     | *                                           |
| 4      | 1      | 1      | 4  | $(a + b, 0)$                 | $(\frac{1}{2}(a - b), -\frac{1}{2}(a - b))$ |
| 5      | 2      | 1      | 5  | $(a, \bar{b})$               | $(0, b)$                                    |
| 6      | 2      | 1      | 6  | $(a, \bar{b})$               | $(0, b)$                                    |
| 7      | 2      | 2      | 7  | $(\bar{a}, b)$               | $(a, 0)$                                    |
| 8      | 2      | 2      | 8  | $(\bar{a}, b)$               | $(a, 0)$                                    |
| 9      | 1      | 2      | 9  | $(a - b, 0)$                 | $(\frac{1}{2}(a + b), \frac{1}{2}(a + b))$  |
| 10     | 3      | 3      | 10 | $(\bar{a}, \bar{b})$         | $(a, b)$                                    |
| 11     | 3      | 3      | 10 | $(\bar{a} + 1, \bar{b} + 1)$ | $(a, b)$                                    |

Here is the block dual to  $SO(4, 1)$ :

|          |   |         |   |   |       |       |   |         |
|----------|---|---------|---|---|-------|-------|---|---------|
| 0( 5,2): | 1 | [C+,rn] | 2 | 0 | (*,*) | (*,*) | 2 | 2       |
| 1( 6,2): | 1 | [C+,rn] | 3 | 1 | (*,*) | (*,*) | 2 | 2       |
| 2( 7,1): | 2 | [C-,i1] | 0 | 3 | (*,*) | (4,*) | 2 | 1,2,1   |
| 3( 8,1): | 2 | [C-,i1] | 1 | 2 | (*,*) | (4,*) | 2 | 1,2,1   |
| 4(10,0): | 3 | [rn,r1] | 4 | 4 | (*,*) | (2,3) | 3 | 2,1,2,1 |

| number | Cartan | length | x  | $\lambda$                | $\nu$    |
|--------|--------|--------|----|--------------------------|----------|
| 0      | 2      | 1      | 7  | $(\bar{b} + 1, a)$       | $(b, 0)$ |
| 1      | 2      | 1      | 8  | $(\bar{b} + 1, a)$       | $(b, 0)$ |
| 2      | 2      | 2      | 7  | $(\bar{a} + 1, b)$       | $(a, 0)$ |
| 3      | 2      | 2      | 8  | $(\bar{a} + 1, b)$       | $(a, 0)$ |
| 4      | 3      | 3      | 10 | $(\bar{a} + 1, \bar{b})$ | $(a, b)$ |

The principal series  $I(x_{10}, (\bar{a}, \bar{b} + 1), (a, b))$  is irreducible, dual to  $SO(5, 0)$ .

## 9 Orientation Numbers for $Sp(4, \mathbb{R})$

See [?, Section 5].

**Definition 9.1** Suppose  $\gamma = (x, \lambda, \nu)$  is a parameter and  $\alpha$  is a nonintegral real root. Define

$$(9.2) \quad t_\alpha(\gamma) = \langle \nu - (\lambda - \rho_r), \alpha^\vee \rangle \pmod{2\mathbb{Z}}$$

Since  $\alpha$  is not integral  $t_\alpha \notin \mathbb{Z}$ ; think of it as an element of  $(0, 1) \cup (1, 2)$ .

We say a real nonintegral root is oriented (with respect to  $\gamma$ ) if  $0 < t_\alpha(\gamma) < 1$ .

**Definition 9.3** Suppose  $\gamma = (x, \lambda, \nu)$  is a parameter. The orientation number  $\ell_0(\gamma)$  is the sum of

the number of pairs of complex nonintegral positive roots  $\{\alpha, -\theta\alpha\}$

and

the number of real nonintegral oriented roots that are positive on  $\gamma$ .

Here are the orientation numbers for  $Sp(4, \mathbb{R})$ .

On Cartan 0 all roots are imaginary and  $\ell_0 = 0$ .

### 9.1 Cartan 1

Consider  $(x, \lambda, \nu) = (x_9, c, x) = (x_9, (c, 0), \frac{1}{2}(x, x))$ , with infinitesimal character  $\gamma = \frac{1}{2}(x + c, x - c)$ . Here  $c \in \mathbb{Z}_{\geq 0}$  and  $x \in \mathbb{R}_{\geq 0}$ . Recall  $(1, 1)$  is the unique positive real root, and  $(2, 0), (0, 2)$  are complex.

The values of the roots are  $\frac{1}{2}(x \pm c), x, c$ . Assume  $\gamma$  is not integral, i.e.  $c, x$  are not integers of the same parity.

**Case 1**

Assume  $x \in \mathbb{Z}$ , but not of the same parity as  $c$ . The integral root system is of type  $D_2 = \{(\pm 1, \pm 1)\}$ . and the nonintegral roots are  $(2, 0), (0, 2)$  (complex),

| Table ??.1 |                               |          |
|------------|-------------------------------|----------|
| $c, x$     | complex pair $(2, 0), (0, 2)$ | $\ell_0$ |
| $c > x$    |                               | 0        |
| $c < x$    | Y                             | 1        |

**Case 2**

Assume  $x \notin \mathbb{Z}$ . The only positive integral root is  $(1, -1)$ , the integral root system is type  $A_1$ , and the nonintegral roots are  $(2, 0), (0, 2)$  (complex) and  $(1, 1)$  (real).

Compute

$$\nu - (\lambda - \rho_r) = \frac{1}{2}(x, x) - (c, 0) + \left(\frac{1}{2}, \frac{1}{2}\right) = \left(-c + \frac{x-1}{2}, \frac{x-1}{2}\right)$$

and

$$t_\alpha = -c + x + 1$$

Therefore  $\alpha$  is oriented if and only if  $1 < x - c < 2 \pmod{2\mathbb{Z}}$ , or equivalently

$$0 < c - x < 1 \pmod{2\mathbb{Z}}$$

So:

| Table ??.2 |                                    |                               |          |          |
|------------|------------------------------------|-------------------------------|----------|----------|
| $c, x$     | range $\pmod{2\mathbb{Z}}$         | complex pair $(2, 0), (0, 2)$ | $(1, 1)$ | $\ell_0$ |
| $c > x$    | $0 < c - x < 1 \pmod{2\mathbb{Z}}$ |                               | Y        | 1        |
|            | $1 < c - x < 2 \pmod{2\mathbb{Z}}$ |                               |          | 0        |
| $c < x$    | $0 < c - x < 1 \pmod{2\mathbb{Z}}$ | Y                             | Y        | 2        |
|            | $1 < c - x < 2 \pmod{2\mathbb{Z}}$ | Y                             |          | 1        |

Note that if we write the infinitesimal character as  $(\gamma_1, \gamma_2)$  then the table becomes



| Table ??.3     |                                                |                               |          |          |
|----------------|------------------------------------------------|-------------------------------|----------|----------|
| $c, x$         | range (mod $2\mathbb{Z}$ )                     | complex pair $(2, 0), (0, 2)$ | $(1, 1)$ | $\ell_0$ |
| $\gamma_2 < 0$ | $0 < \gamma_2 < \frac{1}{2} \pmod{\mathbb{Z}}$ |                               | Y        | 0        |
|                | $\frac{1}{2} < \gamma_2 < 1 \pmod{\mathbb{Z}}$ |                               |          | 1        |
| $\gamma_2 > 0$ | $0 < \gamma_2 < \frac{1}{2} \pmod{\mathbb{Z}}$ | Y                             | Y        | 1        |
|                | $\frac{1}{2} < \gamma_2 < 1 \pmod{\mathbb{Z}}$ | Y                             |          | 2        |

which agrees with [?, ?]

## 9.2 Cartan 2

Consider  $(x, \lambda, \nu) = (x_k, (\bar{a}, b), (x, 0))$  with infinitesimal character  $(x, b)$ . If  $x \in \mathbb{Z}$  this is integral and  $\ell_0 = 0$ , so assume  $x \notin \mathbb{Z}$ .

Compute

$$\nu - (\lambda - \rho_r) = (x, 0) - (\bar{a}, b) + (1, 0) = (x - \bar{a} + 1, -b)$$

and for  $\alpha$  the real root  $(2, 0)$ ,  $t_\alpha = x - \bar{a} + 1$ . So  $\alpha$  is oriented if and only if

$$1 < x - \bar{a} < 2 \pmod{2\mathbb{Z}}$$

| Table ??.1 |           |                            |                           |          |          |
|------------|-----------|----------------------------|---------------------------|----------|----------|
| $x, b$     | $\bar{a}$ | range (mod $2\mathbb{Z}$ ) | complex pair $(1, \pm 1)$ | $(2, 0)$ | $\ell_0$ |
| $x < b$    | 0         | $0 < x < 1$                |                           | Y        | 0        |
| $x < b$    | 0         | $1 < x < 2$                |                           |          | 1        |
| $x < b$    | 1         | $0 < x < 1$                |                           | Y        | 1        |
| $x < b$    | 1         | $1 < x < 2$                |                           |          | 0        |
| $x > b$    | 0         | $0 < x < 1$                | Y                         | Y        | 1        |
| $x > b$    | 0         | $1 < x < 2$                | Y                         |          | 2        |
| $x > b$    | 1         | $0 < x < 1$                | Y                         | Y        | 2        |
| $x > b$    | 1         | $1 < x < 2$                | Y                         |          | 1        |

## 9.3 Cartan 3

Consider  $I(x_{10}, (\bar{a}, \bar{b}), (x, y))$  with  $x > y > 0$ . The nonintegral cases are as follows (not including the case with no integral roots). The third column gives the type of the integral root system, and the last column gives a consequence of the conditions in the first column.

| Case | $x, y$                                           | type  | integral roots |                                        |
|------|--------------------------------------------------|-------|----------------|----------------------------------------|
| 1    | $x \in \mathbb{Z}, y \notin \mathbb{Z}$          | $A_1$ | $(2, 0)$       |                                        |
| 2    | $x \notin \mathbb{Z}, y \in \mathbb{Z}$          | $A_1$ | $(0, 2)$       |                                        |
| 3    | $x + y \in \mathbb{Z}; x - y \notin \mathbb{Z}$  | $A_1$ | $(1, 1)$       | $x, y \notin \mathbb{Z} + \frac{1}{2}$ |
| 4    | $x - y \in \mathbb{Z}; x + y \notin \mathbb{Z}$  | $A_1$ | $(1, -1)$      | $x, y \notin \mathbb{Z} + \frac{1}{2}$ |
| 5    | $x \pm y \in \mathbb{Z}; x, y \notin \mathbb{Z}$ | $D_2$ | $(1, \pm 1)$   | $x, y \in \mathbb{Z} + \frac{1}{2}$    |

We have  $\rho = \rho_r = (2, 1)$  and

$$(9.3.4) \quad \nu - (\lambda - \rho_r) = (x - a + 2, y - b + 1) \equiv (x - a, y - b + 1).$$

**Cartan 3, Case 1:**  $x \in \mathbb{Z}, y \notin \mathbb{Z}$

The nonintegral roots are  $(1, \pm 1)$  and  $(0, 2)$ . They are oriented if:

$$(9.3.5) \quad \begin{aligned} (1, 1) : 1 < x + y - (\bar{a} + \bar{b}) < 2 \pmod{2\mathbb{Z}} \\ (1, -1) : 1 < x - y - (\bar{a} + \bar{b}) < 2 \pmod{2\mathbb{Z}} \\ (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \end{aligned}$$

It is easy to see that exactly one of  $(1, \pm 1)$  is an oriented root.

| Table ??.1 |                            |                        |          |          |
|------------|----------------------------|------------------------|----------|----------|
| $\bar{b}$  | range (mod $2\mathbb{Z}$ ) | $(1, 1)$ xor $(1, -1)$ | $(0, 2)$ | $\ell_0$ |
| 0          | $0 < y < 1$                | Y                      |          | 1        |
|            | $1 < y < 2$                | Y                      | Y        | 2        |
| 1          | $0 < y < 1$                | Y                      | Y        | 2        |
|            | $1 < y < 2$                | Y                      |          | 1        |

**Cartan 3, Case 2:**  $x \notin \mathbb{Z}, y \in \mathbb{Z}$

The nonintegral roots are  $(1, \pm 1)$  and  $(2, 0)$ . Now both  $(1, \pm 1)$  are oriented, or neither are. The condition on  $(2, 0)$  is

$$(9.3.6) \quad (2, 0) : 0 < x - \bar{a} < 1 \pmod{2\mathbb{Z}}$$

| Table ??.2 |                     |                        |                            |                        |          |          |
|------------|---------------------|------------------------|----------------------------|------------------------|----------|----------|
| $\bar{a}$  | $\bar{a} + \bar{b}$ | $x \pmod{2\mathbb{Z}}$ | $x + y \pmod{2\mathbb{Z}}$ | $(1, 1)$ and $(1, -1)$ | $(2, 0)$ | $\ell_0$ |
| 0          | 0                   | $0 < x < 1$            | $0 < x + y < 1$            | Y                      | Y        | 1        |
|            |                     |                        | $1 < x + y < 2$            |                        | Y        | 3        |
|            |                     | $1 < x < 2$            | $0 < x + y < 1$            | Y                      |          | 0        |
|            |                     |                        | $1 < x + y < 2$            |                        |          | 2        |
| 1          | 0                   | $0 < x < 1$            | $0 < x + y < 1$            | Y                      |          | 0        |
|            |                     |                        | $1 < x + y < 2$            |                        |          | 2        |
|            |                     | $1 < x < 2$            | $0 < x + y < 1$            | Y                      | Y        | 1        |
|            |                     |                        | $1 < x + y < 2$            |                        | Y        | 3        |
| 0          | 1                   | $0 < x < 1$            | $0 < x + y < 1$            | Y                      | Y        | 3        |
|            |                     |                        | $1 < x + y < 2$            |                        | Y        | 1        |
|            |                     | $1 < x < 2$            | $0 < x + y < 1$            | Y                      |          | 2        |
|            |                     |                        | $1 < x + y < 2$            |                        |          | 0        |
| 1          | 1                   | $0 < x < 1$            | $0 < x + y < 1$            | Y                      |          | 2        |
|            |                     |                        | $1 < x + y < 2$            |                        |          | 0        |
|            |                     | $1 < x < 2$            | $0 < x + y < 1$            | Y                      | Y        | 3        |
|            |                     |                        | $1 < x + y < 2$            |                        | Y        | 1        |

**Cartan 3, Case 3:**  $x + y \in \mathbb{Z}, x - y \notin \mathbb{Z}$

An example is  $(x, y) = (\frac{3}{4}, \frac{1}{4})$ . The nonintegral roots are  $(2, 0)$ ,  $(0, 2)$  and  $(1, -1)$ . The oriented conditions are:

$$\begin{aligned}
 (2, 0) : 0 < x - \bar{a} < 1 \pmod{2\mathbb{Z}} \\
 (9.3.7)(a) \quad (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \\
 (1, -1) : 1 < (x - \bar{a}) - (y - \bar{b}) < 2 \pmod{2\mathbb{Z}}
 \end{aligned}$$

Recall  $x + y \in \mathbb{Z}$ . For example, suppose  $x + y$  is even, so  $x = -y \pmod{2\mathbb{Z}}$ , so replace  $x$  with  $-y$  everywhere:

$$\begin{aligned}
 (2, 0) : 0 < -y - \bar{a} < 1 \pmod{2\mathbb{Z}} \\
 (9.3.7)(b) \quad (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \\
 (1, -1) : 1 < -2y - \bar{a} + \bar{b} < 2 \pmod{2\mathbb{Z}}.
 \end{aligned}$$

After a few manipulations, including dividing the last line by 2, this is equiv-

alent to

$$\begin{aligned}
 (2, 0) : 1 < y + \bar{a} < 2 \pmod{2\mathbb{Z}} \\
 (9.3.7)(c) \quad (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \\
 (1, -1) : 0 < y + \frac{1}{2}(\bar{a} - \bar{b}) < \frac{1}{2} \pmod{\mathbb{Z}}.
 \end{aligned}$$

If  $x + y$  is odd replace  $x$  with  $-y + 1 \pmod{2\mathbb{Z}}$ , and something similar happens. Here is the answer.

If  $x + y \in 2\mathbb{Z}, x - y \notin \mathbb{Z}$ :

| Table ??.3           |                            |          |          |           |          |
|----------------------|----------------------------|----------|----------|-----------|----------|
| $(\bar{a}, \bar{b})$ | range $\pmod{2\mathbb{Z}}$ | $(2, 0)$ | $(0, 2)$ | $(1, -1)$ | $\ell_0$ |
| $(0, 0)$             | $0 < y < \frac{1}{2}$      |          |          | Y         | 1        |
|                      | $\frac{1}{2} < y < 1$      |          |          |           | 0        |
|                      | $1 < y < \frac{3}{2}$      | Y        | Y        | Y         | 3        |
|                      | $\frac{3}{2} < y < 2$      | Y        | Y        |           | 2        |
| $(1, 0)$             | $0 < y < \frac{1}{2}$      | Y        |          |           | 1        |
|                      | $\frac{1}{2} < y < 1$      | Y        |          | Y         | 2        |
|                      | $1 < y < \frac{3}{2}$      |          | Y        |           | 1        |
|                      | $\frac{3}{2} < y < 2$      |          | Y        | Y         | 2        |
| $(0, 1)$             | $0 < y < \frac{1}{2}$      |          | Y        |           | 1        |
|                      | $\frac{1}{2} < y < 1$      |          | Y        | Y         | 2        |
|                      | $1 < y < \frac{3}{2}$      | Y        |          |           | 1        |
|                      | $\frac{3}{2} < y < 2$      | Y        |          | Y         | 2        |
| $(1, 1)$             | $0 < y < \frac{1}{2}$      | Y        | Y        | Y         | 3        |
|                      | $\frac{1}{2} < y < 1$      | Y        | Y        |           | 2        |
|                      | $1 < y < \frac{3}{2}$      |          |          | Y         | 1        |
|                      | $\frac{3}{2} < y < 2$      |          |          |           | 0        |

If  $x + y \in 2\mathbb{Z} + 1, x - y \notin \mathbb{Z}$ :

| Table ??.4           |                            |          |          |           |          |
|----------------------|----------------------------|----------|----------|-----------|----------|
| $(\bar{a}, \bar{b})$ | range (mod $2\mathbb{Z}$ ) | $(2, 0)$ | $(0, 2)$ | $(1, -1)$ | $\ell_0$ |
| $(0, 0)$             | $0 < y < \frac{1}{2}$      | Y        |          |           | 1        |
|                      | $\frac{1}{2} < y < 1$      | Y        |          | Y         | 2        |
|                      | $1 < y < \frac{3}{2}$      |          | Y        |           | 1        |
|                      | $\frac{3}{2} < y < 2$      |          | Y        | Y         | 2        |
| $(1, 0)$             | $0 < y < \frac{1}{2}$      |          |          | Y         | 1        |
|                      | $\frac{1}{2} < y < 1$      |          |          |           | 0        |
|                      | $1 < y < \frac{3}{2}$      | Y        | Y        | Y         | 3        |
|                      | $\frac{3}{2} < y < 2$      | Y        | Y        |           | 2        |
| $(0, 1)$             | $0 < y < \frac{1}{2}$      | Y        | Y        | Y         | 3        |
|                      | $\frac{1}{2} < y < 1$      | Y        | Y        |           | 2        |
|                      | $1 < y < \frac{3}{2}$      |          |          | Y         | 1        |
|                      | $\frac{3}{2} < y < 2$      |          |          |           | 0        |
| $(1, 1)$             | $0 < y < \frac{1}{2}$      |          | Y        |           | 1        |
|                      | $\frac{1}{2} < y < 1$      |          | Y        | Y         | 2        |
|                      | $1 < y < \frac{3}{2}$      | Y        |          |           | 1        |
|                      | $\frac{3}{2} < y < 2$      | Y        |          | Y         | 2        |

**Cartan 3, Case 4:**  $x + y \notin \mathbb{Z}, x - y \in \mathbb{Z}$

An example is  $(x, y) = (\frac{4}{3}, \frac{1}{3})$ .

The nonintegral roots are  $(2, 0), (0, 2)$  and  $(1, 1)$ . The oriented conditions are

$$\begin{aligned}
 (2, 0) : 0 < x - \bar{a} < 1 \pmod{2\mathbb{Z}} \\
 (9.3.8)(a) \quad (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \\
 (1, 1) : 1 < (x - \bar{a}) + (y - \bar{b}) < 2 \pmod{2\mathbb{Z}}
 \end{aligned}$$

Suppose  $x - y = n \in \mathbb{Z}$ . As in Case 3, (a) becomes

$$\begin{aligned}
 (2, 0) : 0 < y + n - \bar{a} < 1 \pmod{2\mathbb{Z}} \\
 (9.3.8)(b) \quad (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \\
 (1, 1) : \frac{1}{2} < y + \frac{1}{2}(n - \bar{a} - \bar{b}) < 1 \pmod{\mathbb{Z}}.
 \end{aligned}$$

If  $x - y \in 2\mathbb{Z}, x + y \notin \mathbb{Z}$ :

| Table ??.5           |                            |          |          |          |          |
|----------------------|----------------------------|----------|----------|----------|----------|
| $(\bar{a}, \bar{b})$ | range (mod $2\mathbb{Z}$ ) | $(2, 0)$ | $(0, 2)$ | $(1, 1)$ | $\ell_0$ |
| $(0, 0)$             | $0 < y < \frac{1}{2}$      | Y        |          |          | 1        |
|                      | $\frac{1}{2} < y < 1$      | Y        |          | Y        | 2        |
|                      | $1 < y < \frac{3}{2}$      |          | Y        |          | 1        |
|                      | $\frac{3}{2} < y < 2$      |          | Y        | Y        | 2        |
| $(1, 0)$             | $0 < y < \frac{1}{2}$      |          |          | Y        | 1        |
|                      | $\frac{1}{2} < y < 1$      |          |          |          | 0        |
|                      | $1 < y < \frac{3}{2}$      | Y        | Y        | Y        | 3        |
|                      | $\frac{3}{2} < y < 2$      | Y        | Y        |          | 2        |
| $(0, 1)$             | $0 < y < \frac{1}{2}$      | Y        | Y        | Y        | 3        |
|                      | $\frac{1}{2} < y < 1$      | Y        | Y        |          | 2        |
|                      | $1 < y < \frac{3}{2}$      |          |          | Y        | 1        |
|                      | $\frac{3}{2} < y < 2$      |          |          |          | 0        |
| $(1, 1)$             | $0 < y < \frac{1}{2}$      |          | Y        |          | 1        |
|                      | $\frac{1}{2} < y < 1$      |          | Y        | Y        | 2        |
|                      | $1 < y < \frac{3}{2}$      | Y        |          |          | 1        |
|                      | $\frac{3}{2} < y < 2$      | Y        |          | Y        | 2        |

If  $x - y \in 2\mathbb{Z} + 1, x + y \notin \mathbb{Z}$ :

| Table ??.6           |                            |          |          |          |          |
|----------------------|----------------------------|----------|----------|----------|----------|
| $(\bar{a}, \bar{b})$ | range (mod $2\mathbb{Z}$ ) | $(2, 0)$ | $(0, 2)$ | $(1, 1)$ | $\ell_0$ |
| $(0, 0)$             | $0 < y < \frac{1}{2}$      |          |          | Y        | 1        |
|                      | $\frac{1}{2} < y < 1$      |          |          |          | 0        |
|                      | $1 < y < \frac{3}{2}$      | Y        | Y        | Y        | 3        |
|                      | $\frac{3}{2} < y < 2$      | Y        | Y        |          | 2        |
| $(1, 0)$             | $0 < y < \frac{1}{2}$      | Y        |          |          | 1        |
|                      | $\frac{1}{2} < y < 1$      | Y        |          | Y        | 2        |
|                      | $1 < y < \frac{3}{2}$      |          | Y        |          | 1        |
|                      | $\frac{3}{2} < y < 2$      |          | Y        | Y        | 2        |
| $(0, 1)$             | $0 < y < \frac{1}{2}$      |          | Y        |          | 1        |
|                      | $\frac{1}{2} < y < 1$      |          | Y        | Y        | 2        |
|                      | $1 < y < \frac{3}{2}$      | Y        |          |          | 1        |
|                      | $\frac{3}{2} < y < 2$      | Y        |          | Y        | 2        |
| $(1, 1)$             | $0 < y < \frac{1}{2}$      | Y        | Y        | Y        | 3        |
|                      | $\frac{1}{2} < y < 1$      | Y        | Y        |          | 2        |
|                      | $1 < y < \frac{3}{2}$      |          |          | Y        | 1        |
|                      | $\frac{3}{2} < y < 2$      |          |          |          | 0        |

**Cartan 3, Case 5:**  $x \pm y \in \mathbb{Z}, x, y \notin \mathbb{Z}$

In other words  $x, y \in \mathbb{Z} + \frac{1}{2}$ . An example is  $(x, y) = (\frac{3}{2}, \frac{1}{2})$ .

The nonintegral roots are  $(2, 0), (0, 2)$ . The oriented conditions are

$$(9.3.9)(a) \quad \begin{aligned} (2, 0) : 0 < x - \bar{a} < 1 \pmod{2\mathbb{Z}} \\ (0, 2) : 1 < y - \bar{b} < 2 \pmod{2\mathbb{Z}} \end{aligned}$$

$x \pm y \in \mathbb{Z}; x, y \notin \mathbb{Z}$

| Table ???.7          |                        |                        |                  |          |
|----------------------|------------------------|------------------------|------------------|----------|
| $(\bar{a}, \bar{b})$ | $x \pmod{2\mathbb{Z}}$ | $y \pmod{2\mathbb{Z}}$ | oriented         | $\ell_0$ |
| $(\bar{0}, \bar{0})$ | $0 < x < 1$            | $0 < y < 1$            | $(2, 0)$         | 1        |
|                      | $0 < x < 1$            | $1 < y < 2$            | $(2, 0), (0, 2)$ | 2        |
|                      | $1 < x < 2$            | $0 < y < 1$            |                  | 0        |
|                      | $1 < x < 2$            | $1 < y < 2$            | $(0, 2)$         | 1        |
| $(\bar{1}, \bar{0})$ | $0 < x < 1$            | $0 < y < 1$            |                  | 0        |
|                      | $0 < x < 1$            | $1 < y < 2$            | $(0, 2)$         | 1        |
|                      | $1 < x < 2$            | $0 < y < 1$            | $(2, 0)$         | 1        |
|                      | $1 < x < 2$            | $1 < y < 2$            | $(2, 0), (0, 2)$ | 2        |
| $(\bar{0}, \bar{1})$ | $0 < x < 1$            | $0 < y < 1$            | $(2, 0), (0, 2)$ | 2        |
|                      | $0 < x < 1$            | $1 < y < 2$            | $(2, 0)$         | 1        |
|                      | $1 < x < 2$            | $0 < y < 1$            | $(0, 2)$         | 1        |
|                      | $1 < x < 2$            | $1 < y < 2$            |                  | 0        |
| $(\bar{1}, \bar{1})$ | $0 < x < 1$            | $0 < y < 1$            | $(0, 2)$         | 1        |
|                      | $0 < x < 1$            | $1 < y < 2$            |                  | 0        |
|                      | $1 < x < 2$            | $0 < y < 1$            | $(2, 0), (0, 2)$ | 2        |
|                      | $1 < x < 2$            | $1 < y < 2$            | $(2, 0)$         | 1        |

## 10 Composition Series: Integral Infinitesimal Character

Assume the infinitesimal character  $\gamma$  is regular, so  $\gamma = (a, b)$  with  $a > b > 0$ .

**Cartan 1** There are two standard modules  $I(x_9, (a \mp b, 0), \frac{1}{2}(a \pm b, a \pm b))$ ; necessarily  $a \mp b \in \mathbb{Z}$  (depending on which of the two modules we consider).

By the preceding sections the real root  $(1, 1)$  gives reducibility if and only if  $a \in \mathbb{Z}$ , and the complex root  $(2, 0)$  gives reducibility only of  $I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b))$ , again only if  $a \in \mathbb{Z}$ . So we may assume  $a, b \in \mathbb{Z}$ .

First consider  $I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b))$ . Then (cf. (??))

$$I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) = I(x_4, (a + b, 0), \frac{1}{2}(a - b, -a + b));$$

this is the module **#4** in the output of **block**. Using **klbasis** we see this standard module has three composition factors, including the two large discrete series representations at this infinitesimal character.

**Remark 10.1** *Recall that in order to compute the composition factors of the standard modules using **klbasis**, we need to invert the matrix obtained from the values of the KLV polynomials at 1 with the signs coming from the lengths, as in (??).*

So

$$(10.2)(a) \quad \begin{aligned} I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) = \\ J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) \\ + J(x_0, (a, b)) + J(x_1, (a, b)). \end{aligned}$$

Next,  $I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b))$  is block element 9, so **klbasis** says  $J$  has irreducible modules **9, 0, 1, 4, 5, 6** as composition factors (with multiplicity one).

Then:

$$(10.2)(b) \quad \begin{aligned} I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) = \\ J(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) \\ + J(x_0, (a, b)) + J(x_1, (a, b)) \\ + J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) \\ + J(x_7, (\bar{b}, a), (b, 0)) \\ + J(x_8, (\bar{b}, a), (b, 0)) \end{aligned}$$

**Cartan 2**



The standard modules are  $I(x_k, (\epsilon, b), (a, 0))$  and  $I(x_k, (\epsilon, a), (b, 0))$  with  $k = 7, 8$  and  $\epsilon = 0, 1$ . We continue to assume  $a > b > 0$ .

The real root  $(2, 0)$  gives reducibility  $I(x_k, (\bar{a}, b), (a, 0))$  and  $I(x_k, (\bar{b}, a), (b, 0))$  provided  $a, b \in \mathbb{Z}$ . The complex root  $(1, 1)$  contributes to reducibility of  $I(x_k, (\epsilon, b), (a, 0))$ , again only if  $a, b \in \mathbb{Z}$ . So the infinitesimal character is necessarily integral. These representations are in one of two blocks, depending on  $\epsilon$ .

Using **klbasis** we compute the following composition series.

Here are the formulas in the big block.

$I(x_7, (\bar{b}, a), (b, 0))$  (standard module 5):

$$(10.2)(c) \quad I(x_7, (\bar{b}, a), (b, 0)) = J(x_7, (\bar{b}, a), (b, 0)) + J(x_0, (a, b)) + J(x_2, (a, b)).$$

$I(x_8, (\bar{b}, a), (b, 0))$  (standard module 6):

$$(10.2)(d) \quad I(x_8, (\bar{b}, a), (b, 0)) = J(x_8, (\bar{b}, a), (b, 0)) + J(x_1, (a, b)) + J(x_3, (a, b)).$$

$I(x_7, (\bar{a}, b), (a, 0))$  (standard module 7):

$$(10.2)(e) \quad \begin{aligned} I(x_7, (\bar{a}, b), (a, 0)) &= J(x_7, (\bar{a}, b), (a, 0)) + J(x_0, (a, b)) \\ &+ J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) + J(x_7, (\bar{b}, a), (b, 0)) \end{aligned}$$

$I(x_8, (\bar{a}, b), (a, 0))$  (standard module 8):

$$(10.2)(f) \quad \begin{aligned} I(x_8, (\bar{a}, b), (a, 0)) &= J(x_8, (\bar{a}, b), (a, 0)) + J(x_1, (a, b)) \\ &+ J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) + J(x_8, (\bar{b}, a), (b, 0)) \end{aligned}$$

Here are the formulas in the block dual to  $SO(4, 1)$ .

By the preceding  $I(x_k, (\bar{b} + 1, a), (b, 0))$  with  $k = 7, 8$  are both irreducible. These are standard modules **0, 1** from this block.

$I(x_7, (\bar{a} + 1, b), (a, 0))$  (standard module 2):

$$(10.2)(g) \quad \begin{aligned} I(x_7, (\bar{a} + 1, b), (a, 0)) &= J(x_7, (\bar{a} + 1, b), (a, 0)) \\ &+ J(x_7, (\bar{b} + 1, a), (b, 0)) \end{aligned}$$

$I(x_8, (\bar{a} + 1, b), (a, 0))$  (standard module 3):

$$(10.2)(h) \quad \begin{aligned} I(x_8, (\bar{a} + 1, b), (a, 0)) = & J(x_8, (\bar{a} + 1, b), (a, 0)) \\ & + J(x_8, (\bar{b} + 1, a), (b, 0)) \end{aligned}$$

### Cartan 3

These are the modules  $I(x_{10}, (\delta, \epsilon), (a, b))$  with  $\delta, \epsilon = 0, 1$  and  $a > b > 0$ .

First consider the representations in the big block, so  $a, b \in \mathbb{Z}$ .

$I(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b))$  This is block element 11, which has irreducible modules 11, 0, 1, 2, 3, 4, 5, 6, 9, all of multiplicity one.

$$(10.2)(i) \quad \begin{aligned} I(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b)) = & J(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b)) \\ & + J(x_0, (a, b)) + J(x_1, (a, b)) + J(x_2, (a, b)) + J(x_3, (a, b)) \\ & + J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) \\ & + J(x_7, (\bar{b}, a), (b, 0)) + J(x_8, (\bar{b}, a), (b, 0)) \\ & + J(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) \end{aligned}$$

$I(x_{10}, (\bar{a}, \bar{b}), (a, b))$  This is block element 10, which has irreducible modules 10, 0, 1, 4, 5, 6, 7, 8, 9, all of multiplicity one except 4 has multiplicity 2. See Section ??.

$$(10.2)(j) \quad \begin{aligned} I(x_{10}, (\bar{a}, \bar{b}), (a, b)) = & J(x_{10}, (\bar{a}, \bar{b}), (a, b)) \\ & + J(x_0, (a, b)) + J(x_1, (a, b)) \\ & + 2 \times J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) \\ & + J(x_7, (\bar{b}, a), (b, 0)) + J(x_8, (\bar{b}, a), (b, 0)) \\ & + J(x_7, (\bar{a}, b), (a, 0)) + J(x_8, (\bar{a}, b), (a, 0)) \\ & + J(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) \end{aligned}$$

Next, consider the integral block dual to  $SO(4, 1)$ . There is one representation on the split Cartan, number 4.

$$\begin{aligned}
(10.2)(k) \quad I(x_{10}, (\bar{a} + 1, \bar{b}), (a, b)) &= J(x_{10}, (\bar{a} + 1, \bar{b}), (a, b)) \\
&+ J(x_7, (\bar{b} + 1, a), (b, 0)) + J(x_8, (\bar{b} + 1, a), (b, 0)) \\
&+ J(x_7, (\bar{a} + 1, b), (a, 0)) + J(x_8, (\bar{a} + 1, b), (a, 0))
\end{aligned}$$

The principal series  $I(x_{10}, (\bar{a}, \bar{b}+1), (a, b))$  is irreducible (dual to  $SO(5, 0)$ ).

## 11 Character Formulas: Integral Infinitesimal Character

Assume the infinitesimal character  $\gamma$  is regular, so  $\gamma = (a, b)$  with  $a > b > 0$ .

Suppose  $\mu = (x, \lambda, \nu)$ . Use `nblock` to compute the block of  $I(\mu)$ , and `klbasis` to compute  $P_{\mu, \delta}$  for all  $\delta$  in the block. Then:

$$(11.1) \quad J(\mu) = \sum (-1)^{\ell(\mu) - \ell(\delta)} P_{\delta, \mu}(1) I(\delta)$$

Here are character formulas; the numbering is parallel to that of (??)(a-k).

### Cartan 1

$$\begin{aligned}
(11.2)(a) \quad J(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) &= \\
I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) & \\
- I(x_0, (a, b)) - I(x_1, (a, b)). &
\end{aligned}$$

$$\begin{aligned}
(11.2)(b) \quad J(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) &= \\
I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) & \\
+ I(x_0, (a, b)) + I(x_1, (a, b)) + I(x_2, (a, b)) + I(x_3, (a, b)) & \\
- I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) & \\
- I(x_7, (\bar{b}, a), (b, 0)) - I(x_8, (\bar{b}, a), (b, 0)) &
\end{aligned}$$

### Cartan 2

Here are the formulas on the big block.

$$(11.2)(c) \quad J(x_7, (\bar{b}, a), (b, 0)) = I(x_7, (\bar{b}, a), (b, 0)) - I(x_0, (a, b)) - I(x_2, (a, b))$$

$$(11.2)(d) \quad J(x_8, (\bar{b}, a), (b, 0)) = I(x_8, (\bar{b}, a), (b, 0)) - I(x_1, (a, b)) - I(x_3, (a, b))$$

$$(11.2)(e) \quad \begin{aligned} J(x_7, (\bar{a}, b), (a, 0)) &= I(x_7, (\bar{a}, b), (a, 0)) \\ &+ I(x_0, (a, b)) + I(x_1, (a, b)) + I(x_2, (a, b)) \\ &- I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) - I(x_7, (\bar{b}, a), (b, 0)) \end{aligned}$$

$$(11.2)(f) \quad \begin{aligned} J(x_8, (\bar{a}, b), (a, 0)) &= I(x_8, (\bar{a}, b), (a, 0)) \\ &+ I(x_0, (a, b)) + I(x_1, (a, b)) + I(x_3, (a, b)) \\ &- I(x_9, (a + b, 0), \frac{1}{2}(a - b, a - b)) - I(x_8, (\bar{b}, a), (b, 0)) \end{aligned}$$

Here are the formulas on the block dual to  $SO(4, 1)$ :

$$(11.2)(g) \quad J(x_7, (\bar{a} + 1, b), (a, 0)) = I(x_7, (\bar{a} + 1, b), (a, 0)) - I(x_7, (\bar{b} + 1, a), (b, 0))$$

$$(11.2)(h) \quad J(x_8, (\bar{a} + 1, b), (a, 0)) = I(x_8, (\bar{a} + 1, b), (a, 0)) - I(x_8, (\bar{b} + 1, a), (b, 0))$$

### Cartan 3

First consider the big block, with integral infinitesimal character.

$$(11.2)(i) \quad \begin{aligned} J(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b)) &= I(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b)) - I(x_2, (a, b)) - I(x_3, (a, b)) \\ &- I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b)) \end{aligned}$$

See Section ??.

$$\begin{aligned}
(11.2)(j) \quad J(x_{10}, (\bar{a}, \bar{b}), (a, b)) &= I(x_{10}, (\bar{a}, \bar{b}), (a, b)) \\
&\quad - I(x_0, (a, b)) - I(x_1, (a, b)) - I(x_2, (a, b)) - I(x_3, (a, b)) \\
&\quad + I(x_9, (a+b, 0), \frac{1}{2}(a-b, a-b)) \\
&\quad + I(x_7, (\bar{b}, a), (b, 0)) + I(x_8, (\bar{b}, a), (b, 0)) \\
&\quad - I(x_7, (\bar{a}, b), (a, 0)) - I(x_8, (\bar{a}, b), (a, 0)) \\
&\quad - I(x_9, (a-b, 0), \frac{1}{2}(a+b, a+b))
\end{aligned}$$

On the block dual to  $SO(4, 1)$  there is a single formula on the split Cartan:

$$\begin{aligned}
(11.2)(k) \quad J(x_{10}, (\bar{a}+1, \bar{b}), (a, b)) &= I(x_{10}, (\bar{a}+1, \bar{b}), (a, b)) - \\
&\quad J(x_7, (\bar{a}+1, b), (a, 0)) - J(x_8, (\bar{a}+1, b), (a, 0))
\end{aligned}$$

## 12 Composition Series and Character Formulas: Non-Integral Infinitesimal Character

Write  $\gamma = (a, b)$  with  $a > b > 0$ . Let  $\Delta(\gamma)$  be the integral root system:  $\Delta(\gamma) = \{\alpha \mid \langle \gamma, \alpha^\vee \rangle \in \mathbb{Z}\}$ . If this is empty then the standard module is irreducible. Here are the remaining cases.

### 12.1 $\Delta(\gamma) = \{(\pm 1, \pm 1)\}$ (type $D_2$ )

If  $a, b \in \mathbb{Z} + \frac{1}{2}$  then  $\Delta(\gamma) = \{(\pm 1, \pm 1)\}$  is of type  $D_2$ .

On the compact Cartan subgroup or Cartan 2 at least one of the roots  $2e_i$  is imaginary, hence integral, so this case doesn't arise. So we're on the  $\mathbb{C}^*$  or split Cartan.

Recall ((?))  $I(x_9, c, x)$  has infinitesimal character  $\gamma = \frac{1}{2}(x+c, x-c)$ , with  $c \in \mathbb{Z}$ , which gives  $D(\gamma)$  of type  $D_2$  in the following cases:

$$(12.1.1) \quad I(x_9, c, x) \quad (x, c \geq 0, \text{ integers of opposite parity})$$

For example the representations with infinitesimal character  $(\frac{3}{2}, \frac{1}{2})$  are

$$\begin{aligned} I(x_9, 1, 2) &= I(x_9, (1, 0), (1, 1)) \\ I(x_9, 2, 1) &= I(x_9, (2, 0), (\frac{1}{2}, \frac{1}{2})) \end{aligned}$$

These are irreducible:

Suppose  $x > c$ . For example take  $c = 1, x = 2$ , i.e.  $I(x_9, 1, 2) = I(x_9, (1, 0), (1, 1))$ . Using **nblock** (see Section ??) this is  $\lambda - \rho = (-1, -1) = [0, -1]$  and  $\nu = (1, 1) = [0, 1]$ . The infinitesimal character is  $\gamma = [1, \frac{1}{2}] = (\frac{3}{2}, \frac{1}{2})$ .

```
real: nblock
choose Cartan class (one of 0,1,2,3): 1
Choose a KGB element from the following canonical fiber:
  9:  2  [n,C]      9   4   10   *   (0,0)#1 2,1,2
KGB number: 9
rho = [1,1]/1
NEED, on following imaginary coroot, at least given value:
[1,0] (>=-1)
Give lambda-rho: 0 -1
denominator for nu: 1
numerator for nu: 0 1
Name an output file (return for stdout, ? to abandon):
x = 9, gamma = [2,1]/2, lambda = [1,0]/1
Subsystem on dual side is of type A1.A1, with roots 4,6.
Given parameters define element 0 of the following block:
0(0,2):  0  [i2,rn]  0  0   (1,2)  (*,*)  *( 9,  [1,0]= rho+  [0,-1])  1  1
1(1,0):  1  [r2,rn]  2  1   (0,*)  (*,*)  *(10,  [1,0]= rho+  [0,-1])  0  e
2(1,1):  1  [r2,rn]  1  2   (0,*)  (*,*)  *(10,  [1,-1]= rho+  [0,-2])  0  e
KL polynomials (-1)^(l(0)-l(x))*P_{x,0}:
0: 1
```

The KLV polynomial information tells us that this module (module 0) is irreducible.

The two other standard modules here, on the split Cartan subgroup, are  $I(x_{10}, (\bar{1}, \bar{0}), (\frac{3}{2}, \frac{1}{2}))$  and  $I(x_{10}, (\bar{0}, \bar{1}), (\frac{3}{2}, \frac{1}{2}))$ . Using the **nblock** command for these two (principal series) representations, we see that they have composition series

$$\begin{aligned} I(x_{10}, (\bar{1}, \bar{0}), (\frac{3}{2}, \frac{1}{2})) &= J(x_{10}, (\bar{1}, \bar{0}), (\frac{3}{2}, \frac{1}{2})) + J(x_9, 1, 2) \\ I(x_{10}, (\bar{0}, \bar{1}), (\frac{3}{2}, \frac{1}{2})) &= J(x_{10}, (\bar{0}, \bar{1}), (\frac{3}{2}, \frac{1}{2})) + J(x_9, 1, 2) \end{aligned}$$

and character formulas

$$\begin{aligned} J(x_{10}, (\bar{1}, \bar{0}), (\frac{3}{2}, \frac{1}{2})) &= I(x_{10}, (\bar{1}, \bar{0}), (\frac{3}{2}, \frac{1}{2})) - I(x_9, 1, 2) \\ J(x_{10}, (\bar{0}, \bar{1}), (\frac{3}{2}, \frac{1}{2})) &= I(x_{10}, (\bar{0}, \bar{1}), (\frac{3}{2}, \frac{1}{2})) - I(x_9, 1, 2) \end{aligned}$$

More generally for  $x > c > 0$ ,  $x, c$  integers of opposite parity, the standard modules are

$$(12.1.2)(a) \quad \begin{array}{l} I(x_9, c, x) \quad (x > c \geq 0, \text{ integers of opposite parity}) \\ I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, x-c)) \\ I(x_{10}, (\bar{1}, \overline{1+c}), \frac{1}{2}(x+c, x-c)) \end{array}$$

and the character formulas are

$$(12.1.2)(b) \quad \begin{array}{l} J(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, x-c)) = I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, x-c)) - I(x_9, c, x) \\ J(x_{10}, (\bar{1}, \overline{1+c}), \frac{1}{2}(x+c, x-c)) = I(x_{10}, (\bar{1}, \overline{1+c}), \frac{1}{2}(x+c, x-c)) - I(x_9, c, x) \end{array}$$

**Remark 12.1.3** *If we choose  $x > c = 0$  with  $x$  an odd integer, say  $x = 1$ , we get*

```
real: nblock
choose Cartan class (one of 0,1,2,3): 1
Choose a KGB element from the following canonical fiber:
  9:  2  [n,C]    9   4   10   *   (0,0)#1 2,1,2
KGB number: 9
rho = [1,1]/1
NEED, on following imaginary coroot, at least given value:
[1,0] (>=-1)
Give lambda-rho: -1 -1
denominator for nu: 2
numerator for nu: 0 1
Name an output file (return for stdout, ? to abandon):
x = 9, gamma = [0,1]/2, lambda = [0,0]/1
Subsystem on dual side is of type A1.A1, with roots 4,6.
Given parameters define element 0 of the following block:
0(0,2):  0  [i2,rn]  0  0   (1,2)  (*,*)  *( 9,  [0,0]= rho+  [-1,-1])  1
1(1,0):  1  [r2,rn]  2  1   (0,*)  (*,*)  (10,  [0,0]= rho+  [-1,-1])  e
2(1,1):  1  [r2,rn]  1  2   (0,*)  (*,*)  (10,  [0,-1]= rho+  [-1,-2])  e
(cumulated) KL polynomials (-1)^(l(0)-l(x))*P_{x,0}:
0: 1
```

In this case, the block contains only the module  $I(x_9, 0, x)$ ; the parameters for  $I(x_{10}, (\bar{0}, \bar{0}), \frac{1}{2}(x, x))$  and  $I(x_{10}, (\bar{1}, \bar{1}), \frac{1}{2}(x, x))$  are not final; so the irreducible modules  $J(x_{10}, (\bar{0}, \bar{0}), \frac{1}{2}(x, x))$  and  $J(x_{10}, (\bar{1}, \bar{1}), \frac{1}{2}(x, x))$  are zero. The full principal series representations associated to these parameters are actually irreducible, and both equivalent to  $J(x_9, 0, x)$ .

If  $0 \leq x < c$ , we get the same character formulas. For example  $x = 1, c = 2$  gives  $I(x_9, 2, 1) = I(x_9, (2, 0), \frac{1}{2}(1, 1))$  with  $\gamma = (\frac{3}{2}, -\frac{1}{2})$ . In `nblock` coordinates this is  $\lambda = (2, 0)$ ,  $\lambda - \rho = (0, -1) = [1, -1]$ ,  $\nu = (\frac{1}{2}, \frac{1}{2}) = [0, \frac{1}{2}]$ .

```

real: nblock
choose Cartan class (one of 0,1,2,3): 1
Choose a KGB element from the following canonical fiber:
  9:  2  [n,C]      9   4   10   *   (0,0)#1 2,1,2
KGB number: 9
rho = [1,1]/1
NEED, on following imaginary coroot, at least given value:
[1,0] (>=-1)
Give lambda-rho: 1 -1
denominator for nu: 2
numerator for nu: 0 1
Name an output file (return for stdout, ? to abandon):
x = 9, gamma = [4,-1]/2, lambda = [2,0]/1
Subsystem on dual side is of type A1.A1, with roots 4,6.
Given parameters define element 0 of the following block:
0(0,2):  0  [i2,rn]  0  0   (1,2)  (*,*)  *( 9, [2,-2]= rho+  [1,-3])  1
1(1,0):  1  [r2,rn]  2  1   (0,*)  (*,*)  *(10, [2,-2]= rho+  [1,-3])  e
2(1,1):  1  [r2,rn]  1  2   (0,*)  (*,*)  *(10, [2,-1]= rho+  [1,-2])  e
KL polynomials (-1)^(l(0)-l(x))*P_{x,0}:
0: 1

```

The infinitesimal character is not dominant on the coroot  $(0, 1)$ ; however, it is dominant on the integral coroots. The standard modules and character formulas are exactly as in (??) and (??). Using

$$(12.1.4) \quad I(x_{10}(\bar{a}, \bar{b}), (x, y)) \cong I(x_{10}(\bar{a}, \bar{b}), (x, -y)),$$



we write these formulas

$$(12.1.5)(a) \quad \boxed{\begin{array}{l} I(x_9, c, x) \quad (c > 0, x \geq 0, \text{ integers of opposite parity}) \\ I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, -x+c)) \\ I(x_{10}, (\bar{1}, \bar{1+c}), \frac{1}{2}(x+c, -x+c)) \end{array}}$$

and the character formulas

$$(12.1.5)(b) \quad \boxed{\begin{array}{l} J(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, -x+c)) = I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x+c, -x+c)) - I(x_9, c, x) \\ J(x_{10}, (\bar{1}, \bar{1+c}), \frac{1}{2}(x+c, -x+c)) = I(x_{10}, (\bar{1}, \bar{1+c}), \frac{1}{2}(x+c, -x+c)) - I(x_9, c, x) \end{array}}$$

In all remaining cases  $\Delta(\gamma)$  is of type  $A_1$ .

## 12.2 $\Delta(\gamma) = \{\pm(1, -1)\}$ or $\{\pm(1, 1)\}$

As in (??) we are on the  $\mathbb{C}^*$  or split Cartan. On the  $\mathbb{C}^*$  Cartan take

$$I(x_9, c, x) = I(x_9, (c, 0), \frac{1}{2}(x, x)) \quad (x \notin \mathbb{Z}, x \geq 0, c > 0).$$

with

$$\gamma = \frac{1}{2}(x+c, x-c)$$

for which  $(1, -1)$  is the unique positive integral root (and is imaginary). (If  $x < c$  then we can conjugate to the fiber of **x4** to make the infinitesimal character dominant; then  $(1, 1)$  will be the unique positive integral root.)

The standard module is irreducible.

Here is an example; compare Section ??.

```
real: nblock
choose Cartan class (one of 0,1,2,3): 1
Choose a KGB element from the following canonical fiber:
  9:  2  [n,C]    9   4   10   *   (0,0)#1 2,1,2
KGB number: 9
rho = [1,1]/1
NEED, on following imaginary coroot, at least given value:
[1,0] (>=-1)
Give lambda-rho: 0 -1
```

```

denominator for nu: 8
numerator for nu: 0 5
Name an output file (return for stdout, ? to abandon):
x = 9, gamma = [8,1]/8, lambda = [1,0]/1
Subsystem on dual side is of type A1, with roots 4.
Given parameters define element 0 of the following block:
0(0,2):  0  [i2]  0  (1,2)  *( 9,  [1,0]= rho+  [0,-1])  1  1
1(1,0):  1  [r2]  2  (0,*)  *(10,  [1,0]= rho+  [0,-1])  0  e
2(1,1):  1  [r2]  1  (0,*)  *(10, [1,-1]= rho+  [0,-2])  0  e
KL polynomials (-1)~{l(0)-l(x)}*P_{x,0}:
0: 1

```

Recall the situation is  $x \notin \mathbb{Z}, x, c > 0$ . Compare (??) and (??) , which is the same except that  $x \in \mathbb{Z}$  of opposite parity to  $c$ . The standard modules are

$$(12.2.6)(a) \quad \boxed{\begin{aligned} I(x_9, c, x) &= I(x_9, (c, 0), \frac{1}{2}(x, x)) \quad (x \notin \mathbb{Z}, x > c) \\ I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x + c, x - c)) \\ I(x_{10}, (\bar{1}, \bar{1} + c), \frac{1}{2}(x + c, x - c)) \end{aligned}}$$

and the character formulas are as in (??)

(12.2.6)(b)

$$\boxed{\begin{aligned} J(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x + c, x - c)) &= I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x + c, x - c)) - I(x_9, c, x) \\ J(x_{10}, (\bar{1}, \bar{1} + c), \frac{1}{2}(x + c, x - c)) &= I(x_{10}, (\bar{1}, \bar{1} + c), \frac{1}{2}(x + c, x - c)) - I(x_9, c, x) \end{aligned}}$$

As in (??) and (??), if  $x < c$ , it may be convenient to use the formula (??) to write

(12.2.7)

$$\boxed{\begin{aligned} J(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x + c, -x + c)) &= I(x_{10}, (\bar{0}, \bar{c}), \frac{1}{2}(x + c, -x + c)) - I(x_9, c, x) \\ J(x_{10}, (\bar{1}, \bar{1} + c), \frac{1}{2}(x + c, -x + c)) &= I(x_{10}, (\bar{1}, \bar{1} + c), \frac{1}{2}(x + c, -x + c)) - I(x_9, c, x) \end{aligned}}$$

instead.

### 12.3 $\Delta(\gamma) = \{\pm(0, 2)\}$

Consider

$$I(x_k, (\bar{a}, b), (x, 0)) \quad (k = 7, 8; x \notin \mathbb{Z}; x > b)$$

with  $\gamma = (x, b)$  dominant. This standard module is irreducible. On the split Cartan we have

$$(12.3.8) \quad I(x_{10}, (\bar{a}, \bar{b}), (x, b)).$$

The character formula is (recall  $x > b$ ):

(12.3.9)

$$J(x_{10}, (\bar{a}, \bar{b}), (x, b)) = I(x_{10}, (\bar{a}, \bar{b}), (x, b)) - I(x_7, (\bar{a}, b), (x, 0)) - I(x_8, (\bar{a}, b), (x, 0))$$

**Remark 12.3.10** *If we choose  $b = 0$ , nblock yields:*

```

real: nblock
choose Cartan class (one of 0,1,2,3): 2
Choose a KGB element from the following canonical fiber:
  7:  2  [C,n]    5   8      *  10  (0,0)#2 1,2,1
  8:  2  [C,n]    6   7      *  10  (0,1)#2 1,2,1
KGB number: 7
rho = [1,1]/1
NEED, on following imaginary coroot, at least given value:
[0,1] (>=-1)
Give lambda-rho: -1 -1
denominator for nu: 3
numerator for nu: 0 2
Name an output file (return for stdout, ? to abandon):
x = 7, gamma = [2,0]/3, lambda = [0,0]/1
Subsystem on dual side is of type A1, with roots 5.
Given parameters define element 1 of the following block:
0(0,1):  0  [i1]  1   (2,*)  *( 8,  [0,0]= rho+  [-1,-1])  2
1(1,1):  0  [i1]  0   (2,*)  *( 7,  [0,0]= rho+  [-1,-1])  2
2(2,0):  1  [r1]  2   (1,0)  (10,  [0,0]= rho+  [-1,-1])  e
(cumulated) KL polynomials (-1)^(l(1)-l(x))*P_{x,1}:
1: 1

```

*The block contains only the two modules which are related by a cross action through the imaginary root. Once again, the parameter attached to Cartan 3 is not final; the corresponding principal series representation is the direct sum of the two modules 0 and 1.*

## 12.4 $\Delta(\gamma) = \{\pm(2, 0)\}$

Consider

$$I(x_k, (\bar{a}, b), (x, 0)) \quad (k = 7, 8; x \notin \mathbb{Z}; x < b).$$

Now  $\gamma = (x, b)$  is not dominant, but conjugate to  $(b, x)$ ; this is dominant, and gives integral roots  $\pm(2, 0)$ . Probably it is better to think of this as

$$(12.4.11) \quad I(x_\ell, (b, \bar{a}), (0, x)) \quad (\ell = 5, 6; x \notin \mathbb{Z}; x < b).$$

which makes it clear that  $(2, 0)$  is an imaginary root.

These standard modules are irreducible. There is also a standard module on the split Cartan:

$$(12.4.12) \quad I(x_{10}, (\bar{a}, \bar{b}), (x, b))$$

The character formula is precisely as in (??) (recall that now  $x < b$ ):

$$(12.4.13)$$

$$J(x_{10}, (\bar{a}, \bar{b}), (x, b)) = I(x_{10}, (\bar{a}, \bar{b}), (x, b)) - I(x_7, (\bar{a}, b), (x, 0)) - I(x_8, (\bar{a}, b), (x, 0))$$

However it is probably clearer to use (??) (and rule (c) in Section ?? for conjugation by an element of  $W_r$ ) and write this as (for  $x < b$ ):

$$(12.4.14)$$

$$\begin{aligned} & J(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x)) \\ &= I(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x)) - I(x_5, (b, \bar{a}), (0, x)) - I(x_6, (b, \bar{a}), (0, x)) \end{aligned}$$

## 13 c-invariant Hermitian Forms

Every irreducible representation  $J$  has a distinguished c-invariant Hermitian form. We think of this as a virtual  $K$ -representation with coefficients in  $\mathbb{W} = \mathbb{Z}[s]$  where  $s^2 = 1$ . A term  $(p + qs)\mu$  means that the  $K$ -type  $\mu$  has multiplicity  $p + q$ , and occurs  $p$  times with a plus sign, and  $q$  with a minus. In other words the  $\mu$ -isotypic space is  $V_\mu \otimes V[\mu]$  where  $V_\mu$  is the space of  $\mu$  equipped with a positive definite form, and  $V[\mu] = \mathbb{C}^{p+q}$  with a form of signature  $(p, q)$ .

**Definition 13.1** (a)  $J(x, \lambda, \nu)_c$  is the irreducible representation  $J(x, \lambda, \nu)$ , equipped with its canonical  $c$ -invariant Hermitian form, normalized to be positive on the lowest  $K$ -types.

(b) Write  $J_K(x, \lambda, \nu)_c$  for the restriction of  $J(x, \lambda, \nu)$  to  $K$ . I won't always make this distinction.

(c) Suppose  $I_K(x, \lambda)$  is a nonzero standard final limit  $K$ -representation. Let  $I_K(x, \lambda)_c$  denote this module equipped with the canonical  $c$ -invariant Hermitian form which is positive on the (unique) lowest  $K$ -type.

The  $I_K(x, \lambda)_c$  form a basis of the Grothendieck group of  $K$ -representations, equipped with a  $c$ -invariant Hermitian form.

We want to compute formulas of the form

$$(13.2)(a) \quad J_K(x, \lambda, \nu)_c = \sum_{x', \lambda'} a(x', \lambda') I_K(x', \lambda')_c$$

where the sum is over nonzero final standard limit  $K$ -data, and  $a(x', \lambda') \in \mathbb{W}$ . This is an identity in the Grothendieck group of  $K$ -modules with a  $c$ -invariant form.

This proceeds in two steps. We first write  $J(x, \lambda, \nu)_c$  as a linear combination of  $I(x', \lambda', \nu')_c$  with coefficients in  $\mathbb{W}$ . Recall (??) there is a character formula

$$(13.2)(b) \quad J(x, \lambda, \nu) = \sum_{(x', \lambda', \nu')} (-1)^{\ell(x, \lambda, \nu) - \ell(x', \lambda', \nu')} P_{((x', \lambda', \nu'), (x, \lambda, \nu))}(1) I(x', \lambda', \nu')$$

This is an identity in the Grothendieck group of  $(\mathfrak{g}, K)$ -modules and the sum is over the block containing  $J(x, \lambda, \nu)$ .

If  $I(x, \lambda, \nu)$  is a standard module,  $\text{gr} I(x, \lambda, \nu)$  has a distinguished, nondegenerate,  $c$ -invariant form, obtained by deforming  $\nu$  in the outward direction so it becomes irreducible. We denote this  $I(x, \lambda, \nu)_c$ . Note that  $I(x, \lambda, \nu)_c$  is lower semi-continuous in  $\nu$ . As usual write  $I_K(x, \lambda, \nu)_c$  to denote the restriction to  $K$ .

In the equal rank case there is a simple generalization of (b):

$$(13.2)(c) \quad J(x, \lambda, \nu)_c = \sum_{(x', \lambda', \nu')} \epsilon(s) P_{((x', \lambda', \nu'), (x, \lambda, \nu))}(s) I(x', \lambda', \nu')_c$$

where

$$(13.2)(d) \quad \epsilon(s) = s^{\frac{\ell_0(x, \lambda, \nu) - \ell_0(x', \lambda', \nu')}{2}} (-1)^{\ell(x, \lambda, \nu) - \ell(x', \lambda', \nu')}.$$

This is an identity in the Grothendieck group of  $(\mathfrak{g}, K)$ -modules equipped with a  $c$ -invariant form. The sum is over the block containing  $J(x, \lambda, \nu)$ ; taking  $s = 1$  gives (b). The integers  $\ell_0$  are the orientation numbers of Section ??.

The second step is to write  $I_K(x', \lambda', \nu')_c$  in terms of  $I_K(x'', \lambda'')_c$ . This is by *deforming*  $\nu$  to 0, which we defer to Section ??.

Here is how to compute (?). Use `nblock` to define  $J(x, \lambda, \nu)$  and compute its block; this is at possibly nonintegral or singular infinitesimal character  $\gamma$ . Each parameter in the output of `nblock` may have an asterisk, indicating which of the terms are nonzero at  $\gamma$ . The output also includes a computation of the  $P_{\delta, \mu}$  for  $\mu = (x, \lambda, \nu)$ .

Note that the character formula (b) gives the  $c$ -invariant form formula (c) provided  $P_{*,*}$  is constant for all terms occcuring, and all orientation numbers are 0. All orientation numbers are 0 for integral infinitesimal character.

### 13.1 $c$ -Invariant Forms: Integral Infinitesimal Character

Since the orientation numbers are all 0 the character formulas of Section ?? hold as stated unless some  $P_{\mu, \delta}$  is not a constant. The only case in which this happens is formula (??)(i) in which the terms 2,3 have a coefficient of  $q$ .

Therefore formulas (??)(a-h,j,k) are all valid as formulas for the  $c$ -invariant form, except that (??)(i) should read:

$$(13.1.5) \quad \begin{aligned} J(x_{10}, (\bar{a} + 1, \bar{b} + 1), (a, b))_c &= I(x_{10}, (\bar{a}, \bar{b}), (a, b))_c \\ &\quad - sI(x_2, (a, b))_c - sI(x_3, (a, b))_c \\ &\quad - I(x_9, (a - b, 0), \frac{1}{2}(a + b, a + b))_c \end{aligned}$$

## 13.2 c-Invariant Forms: Nonintegral Infinitesimal Character

In the case of nonintegral infinitesimal character, the integral root system is type  $D_2$  or  $A_1$ , and all the  $P$  polynomials are constant. Therefore the only way for a c-invariant form formula to differ from the character formula is from a difference of orientation numbers being odd.

So we have to consider character formulas  $(??)$ ,  $(??)$ ,  $(??)$ ,  $(??)$ , and  $(??)$  and see which of them require a correction due to the orientation numbers.

Cases  $(??)$ ,  $(??)$ :  $I(x_9, c, x)$  with  $x, c$  integers of opposite parity.

Combining  $(??)$  and  $(??)$  gives, for  $x, y \in \mathbb{Z} + \frac{1}{2}$ ,  $x > y > 0$ ,

$$(13.2.6) \quad J(x_{10}, (\bar{a}, \bar{b}), (x, y)) = I(x_{10}, (\bar{a}, \bar{b}), (x, y)) - \begin{cases} I(x_9, x - y, x + y) & x + y = \bar{a} + \bar{b} + 1 \pmod{2\mathbb{Z}} \\ I(x_9, x + y, x - y) & x + y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}} \end{cases}$$

The orientation numbers for  $I(x_{10}, (*, *), \frac{1}{2}(x + c, x - c))$  are given in Table ???.7, and for  $I(x_9, c, x)$  in Table ???.1.

A little monkeying around shows the following.

If  $x + y = \bar{a} + \bar{b} + 1 \pmod{2\mathbb{Z}}$  then

$$\ell_0(x_{10}, (\bar{a}, \bar{b}), (x, y)) = 1, \ell_0(x_9, x - y, x + y) = 1$$

so there is no contribution from the orientation numbers, and  $(??)$  holds as a formula for c-invariant forms as follows.

Assume  $x + y = a + b + 1 \pmod{2\mathbb{Z}}$ , with  $x, y \in \mathbb{Z} + \frac{1}{2}$ ,  $x > y > 0$ . Then (13.2.7)

$$\boxed{J(x_{10}, (\bar{a}, \bar{b}), (x, y))_c = I(x_{10}, (\bar{a}, \bar{b}), (x, y))_c - I(x_9, x - y, x + y)_c.}$$

Now suppose  $x + y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$ . Then

$$\ell_0(x_9, x + y, x - y) = 0.$$

and on the other hand

$$\ell_0(x_{10}, (\bar{a}, \bar{b}), (x, y)) = \begin{cases} 0 & 0 < y - \bar{b} < 1 \\ 2 & 1 < y - \bar{b} < 2 \end{cases}$$

So this gives the first case of a nontrivial orientation number.

So:  $x + y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  implies

$$(13.2.8) \quad \boxed{J(x_{10}, (\bar{a}, \bar{b}), (x, y))_c = I(x_{10}, (\bar{a}, \bar{b}), (x, y))_c - \begin{cases} I(x_9, x + y, x - y)_c & 0 < y - \bar{b} < 1 \\ sI(x_9, x + y, x - y)_c & 1 < y - \bar{b} < 2 \end{cases}}$$

Case (??):  $x \notin \mathbb{Z}$ . This is very similar to cases (??), (??); see (??) and (??).

Suppose  $x - y \in \mathbb{Z}, x + y \notin \mathbb{Z}$ . If  $x - y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  then

$$(13.2.9)(a) \quad J(x_{10}, (\bar{a}, \bar{b}), (x, y)) = I(x_{10}, (\bar{a}, \bar{b}), (x, y)) - I(x_9, x - y, x + y)$$

If  $x - y \neq \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  then  $I(x_{10}, (\bar{a}, \bar{b}), (x, y))$  is irreducible.

Suppose  $x + y \in \mathbb{Z}, x - y \notin \mathbb{Z}$ . If  $x + y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  then

$$(13.2.9)(b) \quad J(x_{10}, (\bar{a}, \bar{b}), (x, y)) = I(x_{10}, (\bar{a}, \bar{b}), (x, y)) - I(x_9, x + y, x - y)$$

If  $x + y \neq \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  then  $I(x_{10}, (\bar{a}, \bar{b}), (x, y))$  is irreducible.

In (a) all terms have  $\ell_0 = 1$  if  $0 < y < \frac{1}{2} \pmod{\mathbb{Z}}$ , and  $\ell_0 = 2$  if  $\frac{1}{2} < y < 1 \pmod{\mathbb{Z}}$ , so this holds as a formula for c-invariant forms. In other words (??) still holds here:  $x - y \in \mathbb{Z}, x + y \notin \mathbb{Z}, x - y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  implies

$$(13.2.10) \quad \boxed{J(x_{10}, (\bar{a}, \bar{b}), (x, y))_c = I(x_{10}, (\bar{a}, \bar{b}), (x, y))_c - I(x_9, x - y, x + y)_c}$$

Similarly (??) still holds here:  $x + y \in \mathbb{Z}, x - y \notin \mathbb{Z}, x + y = \bar{a} + \bar{b} \pmod{2\mathbb{Z}}$  implies

$$(13.2.11) \quad \boxed{J(x_{10}, (\bar{a}, \bar{b}), (x, y))_c = I(x_{10}, (\bar{a}, \bar{b}), (x, y))_c - \begin{cases} I(x_9, x + y, x - y)_c & 0 < y - \bar{b} < 1 \\ sI(x_9, x + y, x - y)_c & 1 < y - \bar{b} < 2 \end{cases}}$$

Case (??):



Now  $x > y = b > 0$  with  $b \in \mathbb{Z}$ . The character formula in this case is:

(13.2.12)

$$J(x_{10}, (\bar{a}, \bar{b}), (x, b)) = I(x_{10}, (\bar{a}, \bar{b}), (x, b)) - I(x_7, (\bar{a}, b), (x, 0)) - I(x_8, (\bar{a}, b), (x, 0))$$

The orientation numbers  $\ell_0$  of all terms are the same, either 1 or 2, so this holds as a formula for c-invariant forms:

(13.2.13)

$$J(x_{10}, (\bar{a}, \bar{b}), (x, b))_c = I(x_{10}, (\bar{a}, \bar{b}), (x, b))_c - I(x_7, (\bar{a}, b), (x, 0))_c - I(x_8, (\bar{a}, b), (x, 0))_c.$$

Case (??):

This is analogous to the last case, with  $x \notin \mathbb{Z}$  and  $b \in \mathbb{Z}$ , except that now  $x < b$ . The character formula is

$$\begin{aligned} J(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x)) \\ = I(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x)) - I(x_7, (\bar{a}, b), (x, 0)) - I(x_8, (\bar{a}, b), (x, 0)). \end{aligned}$$

If  $1 < x - a < 2 \pmod{2\mathbb{Z}}$  then the orientation numbers of all terms are  $\ell_0 = 1$ ; if  $0 < x - a < 1 \pmod{2\mathbb{Z}}$  then  $I(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x))$  has orientation number 2, while the other terms have  $\ell_0 = 0$ . So we have

(13.2.14)

$$\begin{aligned} J(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x))_c &= I(x_{10}, (\overline{b+1}, \overline{a+1}), (b, x))_c \\ &\quad \begin{cases} -sI(x_7, (\bar{a}, b), (x, 0))_c - sI(x_8, (\bar{a}, b), (x, 0))_c & 0 < x - \bar{a} < 1 \\ -I(x_7, (\bar{a}, b), (x, 0))_c - I(x_8, (\bar{a}, b), (x, 0))_c & 1 < x - \bar{a} < 2 \end{cases} \end{aligned}$$

## 14 Deforming $\nu$ to 0

In the previous section we reduced the computation of  $J(x, \lambda, \nu)_c$  to computing  $I(x', \lambda', \nu')_c$  (see (??)). In this section we discuss how to write

$$(14.1) \quad I(x, \lambda, \nu)_c = \sum_{x', \lambda'} b(x', \lambda') I_K(x', \lambda')_c$$

for  $b(x', \lambda') \in \mathbb{W}$ , and the sum is over nonzero final standard limit  $K$ -data. This proceeds by deformation to  $\nu = 0$ , and by induction, which requires using (??) along the way.

Fix  $(x, \lambda, \nu)$  and consider  $I(x, \lambda) = I(x, \lambda, 0)$ . Assume for the moment that  $I(x, \lambda)$  is final (and nonzero), so  $I_K(x, \lambda)$  is a final limit standard  $K$ -representation.

After deforming  $\nu$  if necessary, we can assume  $I(x, \lambda, t_k \nu)$  is reducible for finitely many  $0 < t_1 < \dots < t_n \leq 1$ . This implies

$$I(x, \lambda, t\nu)_c = I_K(x, \lambda)_c \quad (t < t_1)$$

Assume we've computed  $I(x, \lambda, t\nu)_c$  for  $t_{k-1} \leq t < t_k$ .

Write  $\gamma = (x, \lambda, t_k \nu)$ .

First compute the composition factors  $J(\gamma')$  of  $I(\gamma)$ , and the polynomials  $Q(\gamma', \gamma)$ . (Recall these are the polynomials satisfying  $I(\gamma) = \sum_{\gamma'} Q(\gamma', \gamma)(1)J(\gamma')$ . Currently `nblock` computes  $P(\gamma', \gamma)$ . See Section ??.)

The  $c$ -invariant form changes sign on the odd levels of the Jantzen filtration. What this amounts to is the following.

For each  $\gamma'$  with  $Q(\gamma', \gamma) \neq 0$  write

$$(14.10)(a) \quad Q(\gamma', \gamma)(q) = \sum_{n=0}^{\ell(\gamma)-\ell(\gamma')} a_n(\gamma', \gamma) q^{\frac{\ell(\gamma)-\ell(\gamma')-n}{2}}$$

Note that  $a_n(\gamma', \gamma) = 0$  unless  $\ell(\gamma) - \ell(\gamma') = n \pmod{2\mathbb{Z}}$ , so

$$(14.10)(b) \quad Q(\gamma', \gamma)(q) = \sum_{\substack{n=0 \\ n \equiv \ell(\gamma)-\ell(\gamma')}}^{\ell(\gamma)-\ell(\gamma')} a_n(\gamma', \gamma) q^{\frac{\ell(\gamma)-\ell(\gamma')-n}{2}}$$

Then  $a_n$  is the multiplicity of  $J(\gamma')$  in level  $n$  of the Jantzen filtration. Note that  $\gamma'$  can occur in an odd level of the Jantzen filtration of  $I(\gamma)$  only if  $\ell(\gamma) - \ell(\gamma')$  is odd. Therefore

$$(14.11) \quad \begin{aligned} I(x, \lambda, t_k \nu)_c &= I(x, \lambda, t_{k-1} \nu)_c \\ &+ (1-s) \sum_{\substack{\gamma' \\ \ell(\gamma')-\ell(\gamma) \text{ odd}}} \sum_{n \text{ odd}} s^{\frac{\ell(\gamma)-\ell(\gamma')-n}{2}} s^{\frac{\ell_0(\gamma)-\ell_0(\gamma')}{2}} a_n(\gamma', \gamma) J(\gamma')_c \end{aligned}$$

By (??)(b) the inner sum, (after pulling out the  $\ell_0$  term), is just  $Q(\gamma', \gamma)(s)$ . So:

$$(14.12) \quad \boxed{\begin{aligned} I(x, \lambda, t_k \nu)_c &= I(x, \lambda, t_{k-1} \nu)_c \\ &+ (1-s) \sum_{\substack{\gamma' \\ \ell(\gamma')-\ell(\gamma) \text{ odd}}} s^{\frac{\ell_0(\gamma)-\ell_0(\gamma')}{2}} Q(\gamma', \gamma)(s) J(\gamma')_c \end{aligned}}$$

Here is the big block for  $Sp(4, \mathbb{R})$ .

### length 0

Of course there is nothing to do here:

$$(14.13) \quad J(x_k, (a, b))_c = I_K(x_k, (a, b))_c \quad (0 \leq k \leq 3)$$

### length 1, Cartan 1

Consider  $I(x_9, (a+b, 0), \frac{1}{2}(a-b, a-b))$ . It is easier to use the other coordinates  $I(x_9, c, x) = I(x_9, (c, 0), \frac{1}{2}(x, x))$  (??). Notice that length 1 implies  $c > x$ .

The real root  $(1, 1)$  gives reducibility if and only if  $x = c \pmod{2}$ . The complex root  $(2, 0)$  gives reducibility if and only if  $x = c \pmod{2}$  and  $x \geq c$ .

Here is how to pass back and forth. If  $c > x$ :

$$I(x_9, c, x) = I(x_9, (a+b, 0), \frac{1}{2}(a-b, a-b)) \quad (a = \frac{1}{2}(c+x), b = \frac{1}{2}(c-x))$$

and these have length 1, while if  $c \leq x$ :

$$I(x_9, c, x) = I(x_9, (a-b, 0), \frac{1}{2}(a+b, a+b)) \quad (a = \frac{1}{2}(c+x), b = \frac{1}{2}(x-c))$$

of length 2.

Suppose  $c$  is even. Then

$$(14.14)(a) \quad I(x_9, c, x)_c = I_K(x_9, c)_c \quad (0 \leq x < 2)$$

(This module is not final, so we can write it as a sum of two limits of discrete series...but ignore this for this calculation.) If  $c > 2$  then, taking  $x = 2$ , (??)(a) gives

$$(14.14)(b) \quad I(x_9, c, 2) = J(x_9, c, 2) + J(x_0, \frac{1}{2}(c+2, c-2)) + J(x_1, \frac{1}{2}(c+2, c-2))$$

Apply (??) (or (??)). Since the reducibility points are at integral infinitesimal character the orientation numbers are 0, and for  $2 \leq x < 4$ :

(14.14)(c)

$$\begin{aligned} I(x_9, c, x)_c &= I_K(x_9, c)_c \\ &+ (1-s)[I_K(x_0, \frac{1}{2}(c+2, c-2))_c + I_K(x_1, \frac{1}{2}(c+2, c-2))_c] \end{aligned}$$

Similarly if  $c > 4$  then, taking  $x = 4$ , (??)(a) gives

$$(14.14)(d) \quad I(x_9, c, 4) = J(x_9, c, 4) + J(x_0, \frac{1}{2}(c+4, c-4)) + J(x_1, \frac{1}{2}(c+4, c-4))$$

and by (??) for  $(4 \leq x < 6)$ :

(14.14)(e)

$$\begin{aligned} \mathcal{I}(x_9, c, x)_c &= I_K(x_9, c)_c \\ &\quad + (1-s)[I_K(x_0, \tfrac{1}{2}(c+2, c-2))_c + I_K(x_1, \tfrac{1}{2}(c+2, c-2))_c] \\ &\quad + (1-s)[I_K(x_0, \tfrac{1}{2}(c+4, c-4))_c + I_K(x_1, \tfrac{1}{2}(c+4, c-4))_c] \end{aligned}$$

By induction on  $x$  we see for  $c$  even,  $x < c$ :

(14.14)(f)

$$\begin{aligned} I(x_9, c, x)_c &= I_K(x_9, c)_c \\ &\quad + (1-s) \sum_{k=1}^{\lfloor \frac{x}{2} \rfloor} [I_K(x_0, \tfrac{1}{2}(c+2k, c-2k))_c + I_K(x_1, \tfrac{1}{2}(c+2k, c-2k))_c] \end{aligned}$$

Similarly for  $c$  odd,  $x < c$ :

$$(14.14)(g) \quad \begin{aligned} I(x_9, c, x)_c &= I_K(x_9, c)_c \\ &\quad + (1-s) \sum_{k=1}^{\frac{x+1}{2}} [I_K(x_0, \tfrac{1}{2}(c+(2k-1), c-(2k-1)))_c \\ &\quad + I_K(x_1, \tfrac{1}{2}(c+(2k-1), c-(2k-1)))_c] \end{aligned}$$

In particular at  $\rho$  take  $c = 3, x = 1$ :

(14.14)(h)

$$I(x_9, 3, 1)_c = I_K(x_9, 3)_c + (1-s)[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c]$$

## length 1, Cartan 2

Consider the representations

$$I(x_k, (\delta, a), (x, 0)) \quad (k = 7, 8; a \in \mathbb{Z}_{\geq 0})$$

which have length 1 if  $x < a$ . Such a representation is reducible if and only if  $x = \delta \pmod{2}$ .

Take  $k = 7$  and  $\delta = 0$ . There will be reducibility at  $x = 0, 2, \dots$ . We start with

$$(14.15)(a) \quad I(x_7, (\bar{0}, a), (0, 0)) = I_K(x_7, (\bar{0}, a)).$$

For  $0 \leq x < 2 \leq a$ , we get

$$I(x_7, (\bar{0}, a), (x, 0))_c = I_K(x_7, (\bar{0}, a))_c.$$

If  $2 < a$  then taking  $x = 2$  in (??)(c) gives

$$\begin{aligned} I(x_7, (\bar{0}, a), (2, 0)) &= J(x_7, (\bar{0}, a), (2, 0)) \\ &\quad + J(x_0, (a, 2)) + J(x_2, (a, 2)) \end{aligned}$$

so by (??), if  $(2 \leq x < 4)$ :

$$\begin{aligned} I(x_7, (\bar{0}, a), (x, 0))_c &= I_K(x_7, (\bar{0}, a), (x, 0))_c \\ &\quad + (1 - s)[J(x_0, (a, 2)) + J(x_2, (a, 2))_c] \end{aligned}$$

Repeating this as in the previous case we conclude, for  $x < a$ :

$$(14.15)(b) \quad \boxed{\begin{aligned} I(x_7, (\bar{0}, a), (x, 0))_c &= I_K(x_7, (\bar{0}, a))_c \\ &\quad + (1 - s) \sum_{k=1}^{\lfloor \frac{x}{2} \rfloor} [I_K(x_0, (a, 2k))_c + I_K(x_2, (a, 2k))_c] \end{aligned}}$$

Similarly using (??)(d)

$$(14.15)(c) \quad \boxed{\begin{aligned} I(x_8, (\bar{0}, a), (x, 0))_c &= I_K(x_8, (\bar{0}, a))_c \\ &\quad + (1 - s) \sum_{k=1}^{\lfloor \frac{x}{2} \rfloor} [I_K(x_1, (a, 2k))_c + I_K(x_3, (a, 2k))_c] \end{aligned}}$$

The cases  $k = 7, 8$ ,  $\delta = \bar{1}$  are similar. The results are:

$$(14.15)(d) \quad \boxed{\begin{aligned} I(x_7, (\bar{1}, a), (x, 0))_c &= I_K(x_7, (\bar{1}, a))_c \\ &\quad + (1 - s) \sum_{k=1}^{\lfloor \frac{x+1}{2} \rfloor} [I_K(x_0, (a, 2k - 1))_c + I_K(x_2, (a, 2k - 1))_c] \end{aligned}}$$

$$(14.15)(e) \quad \boxed{\begin{aligned} I(x_8, (\bar{1}, a), (x, 0))_c &= I_K(x_8, (\bar{1}, a))_c \\ &\quad + (1 - s) \sum_{k=1}^{\lfloor \frac{x+1}{2} \rfloor} [I_K(x_1, (a, 2k - 1))_c + I_K(x_3, (a, 2k - 1))_c] \end{aligned}}$$

Consider the case of infinitesimal character  $\rho = (2, 1)$ . Formulas (??)(b-e) specialize as follows.

Take  $a = 2, x = 1$  in (??)(b,c):

$$(14.16)(a) \quad I(x_7, (\bar{0}, 2), (1, 0))_c = I_K(x_7, (\bar{0}, 2))_c$$

$$(14.16)(b) \quad I(x_8, (\bar{0}, 2), (1, 0))_c = I_K(x_8, (\bar{0}, 2))_c$$

Take  $a = 2, x = 1$  in (??)(d,e):

$$(14.16)(c) \quad \boxed{I(x_7, (\bar{1}, 2), (1, 0))_c = I_K(x_7, (\bar{1}, 2))_c + (1 - s)[J(x_0, (2, 1))_c + J(x_2, (2, 1))_c]}$$

$$(14.16)(d) \quad \boxed{I(x_8, (\bar{1}, 2), (1, 0))_c = I_K(x_8, (\bar{1}, 2))_c + (1 - s)[J(x_1, (2, 1))_c + J(x_3, (2, 1))_c]}$$

This completes the length 1 portion of our program. From now on we'll only include some special cases of  $a$  and  $b$ .

### length 2, Cartan 1

These are the modules  $I(x_9, c, x) = I(x_9, (c, 0), \frac{1}{2}(x, x))$  with  $c \leq x$ . This is reducible due to the real root  $(1, 1)$  if  $x = c \pmod{2}$ . It is reducible due to the complex root  $(2, 0)$  provided this is integral, which is again the condition  $x = c \pmod{2}$ . So if  $c$  is odd this is reducible at  $x = 1, 3, \dots, c, c + 2, \dots$

Take  $c = 1$ . Then  $x = 3$  gives infinitesimal character  $(\frac{x+c}{2}, \frac{x-c}{2}) = (2, 1) = \rho$ .

Starting at  $x = 0$ :

$$(14.17)(a) \quad I(x_9, 1, x) = I_K(x_9, 1) \quad (x < 1).$$

Reducibility at  $x = 1$ :

$$(14.17)(b) \quad I(x_9, 1, 1) = J(x_9, 1, 1) + J(x_0, (1, 0)) + J(x_1, (1, 0))$$

The last two terms are limits of large discrete series. By (??) (since we're at integral infinitesimal character all orientation numbers are 0):

$$(14.17)(c) \quad I(x_9, 1, 1)_c = I_K(x_9, 1)_c + (1 - s)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c]$$

Reducibility at  $x = 3$ ; the infinitesimal character is  $\frac{1}{2}(3+1, 3-1) = (2, 1) = \rho$ , so this is directly from (??)(b):

$$\begin{aligned}
 (14.17)(d) \quad I(x_9, 1, 3) &= J(x_9, 1, 3)^2 \\
 &\quad + J(x_0, (2, 1))^0 + J(x_1, (2, 1))^0 \\
 &\quad + J(x_9, 3, 1)^1 \\
 &\quad + J(x_7, (\bar{1}, 2), (1, 0))^1 + J(x_8, (\bar{1}, 2), (1, 0))^1
 \end{aligned}$$

with lengths denoted by superscripts. Therefore (no orientation numbers here; and recall that only modules with odd length difference occur in the sum (??)):

$$\begin{aligned}
 (14.17)(e) \quad I(x_9, 1, 3)_c &= I(x_9, 1, 1)_c \\
 &\quad + (1-s)[J(x_9, 3, 1)_c + J(x_7, (\bar{1}, 2), (1, 0))_c + J(x_8, (\bar{1}, 2), (1, 0))_c]
 \end{aligned}$$

We'll plug in (c) for the first term.

Now for the first time we need to express the c-invariant form on an irreducible (each of the three in the last line) in terms of c-invariant forms on standards, using Section ??, which goes back to the character formulas of Section ?? in this case. Thus by (??)(a)

$$(14.17)(f) \quad J(x_9, 3, 1)_c = I(x_9, 3, 1)_c - J(x_0, (2, 1))_c - J(x_1, (2, 1))_c$$

and plug in (??) for  $I(x_9, 3, 1)$  to give:

$$\begin{aligned}
 (14.17)(g) \quad J(x_9, 3, 1)_c &= I_K(x_9, 3)_c \\
 &\quad + (1-s)[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c] \\
 &\quad - I_K(x_0, (2, 1))_c - I_K(x_1, (2, 1))_c
 \end{aligned}$$

which simplifies to

$$(14.17)(h) \quad \boxed{J(x_9, 3, 1)_c = I_K(x_9, 3)_c - s[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c]}$$

Similarly (??)(c) says:

$$\begin{aligned}
 (14.17)(i) \quad I(x_7, (\bar{1}, 2), (1, 0))_c &= I(x_7, (\bar{1}, 2), (1, 0))_c - I(x_0, (2, 1))_c - I(x_2, (2, 1))_c.
 \end{aligned}$$

Use (??) to expand

$$\begin{aligned}
 (14.17)(j) \quad I(x_7, (\bar{1}, 2), (1, 0))_c &= I_K(x_7, (\bar{1}, 2))_c + (1-s)[J(x_0, (2, 1))_c + J(x_2, (2, 1))_c]
 \end{aligned}$$

and so

(14.17)(k)

$$J(x_7, (\bar{1}, 2), (1, 0))_c = I_K(x_7, (\bar{1}, 2))_c - s[I_K(x_0, (2, 1))_c + I_K(x_2, (2, 1))_c]$$

Finally (??)(d) says:

(14.17)(l)

$$J(x_8, (\bar{1}, 2), (1, 0))_c = I(x_8, (\bar{1}, 2), (1, 0))_c - I(x_1, (2, 1))_c - I(x_3, (2, 1))_c.$$

and using (??) to expand

(14.17)(m)

$$I(x_8, (\bar{1}, 2), (1, 0))_c = I_K(x_8, (\bar{1}, 2))_c + (1 - s)[J(x_1, (2, 1))_c + J(x_3, (2, 1))_c]$$

gives

(14.17)(n)

$$J(x_8, (\bar{1}, 2), (1, 0))_c = I_K(x_8, (\bar{1}, 2))_c - s[I_K(x_1, (2, 1))_c + I_K(x_3, (2, 1))_c].$$

Plugging (c), (h), (k) and (n) into (e) gives:

(14.17)(o)

$$\begin{aligned} I(x_9, 1, 3)_c &= I_K(x_9, 1)_c + (1 - s)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c] \\ &\quad + (1 - s)\{I_K(x_9, 3)_c - s[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c]\} \\ &\quad + (1 - s)\{I_K(x_7, (\bar{1}, 2))_c - s[I_K(x_0, (2, 1))_c + I_K(x_2, (2, 1))_c]\} \\ &\quad + (1 - s)\{I_K(x_8, (\bar{1}, 2))_c - s[I_K(x_1, (2, 1))_c + I_K(x_3, (2, 1))_c]\}. \end{aligned}$$

Grouping terms finally gives:

(14.17)(p)

$$\begin{aligned} I(x_9, 1, 3)_c &= I_K(x_9, 1)_c + (1 - s)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c] \\ &\quad + (1 - s)\{2 \times I_K(x_0, (2, 1))_c + 2 \times I_K(x_1, (2, 1))_c + I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c\} \\ &\quad + (1 - s)\{I_K(x_9, 3)_c + I_K(x_7, \bar{1}, 2))_c + I_K(x_8, \bar{1}, 2))_c\}. \end{aligned}$$

**length 2, Cartan 2** We compute  $I(x_7, (\bar{2}, 1), (2, 0))_c$  and  $I(x_8, (\bar{2}, 1), (2, 0))_c$ .

For  $k = 7, 8$ , let's deform  $I(x_k, (\bar{2}, 1), (2, 0))$  to  $I(x_k, (\bar{2}, 1), (0, 0))$ . By the parity condition  $I(x_k, (\bar{2}, 1), (x, 0))$  is reducible only if  $x = \bar{2} \pmod{2\mathbb{Z}}$ . We're taking  $0 \leq x \leq 2$ , so this occurs only at the endpoints  $x = 0, x = 2$ .

At  $x = 0$  this has to do with the fact that  $I(x_k, (\bar{2}, 1), (0, 0))$  is not final. Ignoring this for the moment, consider  $I(x_k, (\bar{2}, 1), (2, 0))$ , at infinitesimal character  $(2, 1)$ .



The composition series at  $(2, 1)$  are given by (??)(e,f) (with lengths given by superscripts)

$$\begin{aligned}
 (14.18)(a) \quad I(x_7, (\bar{2}, 1), (2, 0))^2 &= J(x_7, (\bar{2}, 1), (2, 0))^2 + J(x_0, (2, 1))^0 \\
 &\quad + J(x_9, 3, 1)^1 + J(x_7, (\bar{1}, 2), (1, 0))^1 \\
 I(x_8, (\bar{2}, 1), (2, 0)) &= J(x_8, (\bar{2}, 1), (2, 0))^2 + J(x_1, (2, 1))^0 \\
 &\quad + J(x_9, 3, 1)^1 + J(x_8, (\bar{1}, 2), (1, 0))^1
 \end{aligned}$$

Considering terms of odd length, with the orientation numbers being all 0, (??) gives

$$\begin{aligned}
 (14.18)(b) \quad I(x_7, (\bar{2}, 1), (2, 0))_c &= I(x_7, (\bar{2}, 1), (0, 0))_c \\
 &\quad + (1-s)[J(x_9, 3, 1)_c + J(x_7, (\bar{1}, 2), (1, 0))_c] \\
 I(x_8, (\bar{2}, 1), (2, 0))_c &= I(x_8, (\bar{2}, 1), (0, 0))_c \\
 &\quad + (1-s)[J(x_9, 3, 1)_c + J(x_8, (\bar{1}, 2), (1, 0))_c]
 \end{aligned}$$

Here we are using that  $I(x_k, (\bar{2}, 1), (2-\epsilon, 0))_c = I(x_k, (\bar{2}, 1), (0, 0))_c$ , since there is no reducibility for  $0 < x < 2$ . We know  $J(x_9, 3, 1)_c$ ,  $J(x_7, (\bar{1}, 2), (1, 0))_c$ ,  $J(x_8, (\bar{1}, 2), (1, 0))_c$  from (??)(h),(k) and (n), respectively. Also use (??) to eliminate the terms  $I(x_k, (\bar{2}, 1), (0, 0))_c$  with  $k = 7, 8$ . Plugging these in gives:

$$\begin{aligned}
 (14.18)(c) \quad I(x_7, (\bar{2}, 1), (2, 0))_c &= I(x_0, (1, 0))_c + I(x_2, (1, 0))_c \\
 &\quad + (1-s)[\{I_K(x_9, 3)_c - s[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c]\} \\
 &\quad + \{I_K(x_7, (\bar{1}, 2))_c - s[I_K(x_0, (2, 1)) + I_K(x_2, (2, 1))]\}]
 \end{aligned}$$

and this can be rewritten (recall that  $(1-s)(-s) = 1-s$ )

$$\begin{aligned}
 (14.18)(d) \quad & \boxed{I(x_7, (\bar{2}, 1), (2, 0))_c = I(x_0, (1, 0))_c + I(x_2, (1, 0))_c} \\
 & \quad + (1-s)[I_K(x_9, 3)_c + I_K(x_7, (\bar{1}, 2))_c \\
 & \quad + 2 \times I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_2, (2, 1))_c].
 \end{aligned}$$

Similarly

$$\begin{aligned}
 (14.18)(e) \quad & \boxed{I(x_8, (\bar{2}, 1), (2, 0))_c = I(x_1, (1, 0))_c + I(x_3, (1, 0))_c} \\
 & \quad + (1-s)[I_K(x_9, 3)_c + I_K(x_8, (\bar{1}, 2))_c \\
 & \quad + I(x_0, (2, 1))_c + 2 \times I(x_1, (2, 1))_c + I(x_3, (2, 1))_c].
 \end{aligned}$$

## 15 Digression: computing the $Q$ polynomials

In the setting of the `atlas` software, suppose we have a block, with elements `0-n`. The `klbasis` command gives the polynomials  $P_{\gamma,\gamma'}(q)$  on the block.

For example consider this part of the output for the big block of  $Sp(4, \mathbb{R})$ :

```
10:  0:  1
      1:  1
      2:  1
      3:  1
      4:  1
      5:  1
      6:  1
      7:  1
      8:  1
      9:  1
     10:  1
```

This says the polynomials  $P_{0,10}(q), P_{1,10}(q), \dots, P_{10,10}(q)$  are all 1. Recall this means

$$J(10) = \sum_k (-1)^{\ell(10) - \ell(k)} P_{k,10}(q) I(k).$$

Taking lengths into account (from the `block` command) gives

$$J(10) = I(10) - I(9) - I(8) - I(7) + I(6) + I(5) + I(4) - I(3) - I(2) - I(1) - I(0)$$

This is the character formula for the trivial representation. See (??)(j).

On the other hand we're interested in the  $Q$  polynomials, which satisfy

$$I(n) = \sum_k Q_{k,n}(1) J(k)$$

Here's how to get these from `klbasis`.

Make  $G$  the dual group, and  $G^\vee$  the group. Then

$$Q_{k,n}(q) = P_{n^\vee, k^\vee}(q)$$

where  $k \rightarrow k^\vee$  is the duality map, given by `dualmap`.

For example take  $G = SO(3, 2)$ ,  $G^\vee = Sp(4, \mathbb{R})$ . Consider this part of the output of `klbasis`:

7: 0: 2  
1: 1  
2: 1  
3: 1  
4: 1  
7: 1

Also `dualmap` gives

[10, 11, 9, 7, 8, 5, 6, 4, 0, 1, 2, 3]

The first line says  $P_{0,7}(q) = 2$  for  $SO(3, 2)$ , and `dualmap` says  $0^\vee = 10, 7^\vee = 4$ , so

$$2 = P_{0,7}^{SO}(q) = Q_{4,10}^{Sp}(q),$$

which says  $J(4)$  has multiplicity 2 in  $I(10)$  for  $Sp(4, \mathbb{R})$ . See (??)(j).

## 16 Second Digression: An alternative version of the calculation

Recall we used (??) to write the form on a standard module  $I(x, \lambda, \nu)$  in terms of forms on a standard module with smaller  $\nu$  parameter, and forms on more tempered irreducible modules  $J(\gamma')$ :

$$(16.1) \quad I(x, \lambda, t_k \nu)_c = I(x, \lambda, t_{k-1} \nu)_c + (1-s) \sum_{\substack{\gamma' \\ \ell(\gamma') - \ell(\gamma) \text{ odd}}} s^{\frac{\ell_0(\gamma) - \ell_0(\gamma')}{2}} Q_{\gamma', \gamma}(s) J(\gamma')_c$$

We proceeded by induction, assuming we'd already computed the terms  $J(\gamma')_c$ .

Here is an equivalent formulation, which combines the two steps, and is easier from a computational point of view. We restate this in a self-contained, single step.

### Inductive algorithm

Suppose we are given  $I(\gamma) = I(x, \lambda, \nu)$ . Typically this is at singular, and possibly nonintegral, infinitesimal character. For small  $\epsilon > 0$ ,  $I(x, \lambda, (1+\epsilon)\nu)$

is irreducible. By (??) we have

$$(16.2)(a) \quad \begin{aligned} I(\gamma)_c &= I(x, \lambda, (1 - \epsilon)\nu)_c \\ &+ (1 - s) \sum_{\substack{\gamma' < \gamma \\ \ell(\gamma) - \ell(\gamma') \text{ odd}}} s^{\frac{\ell_0(\gamma) - \ell_0(\gamma')}{2}} Q_{\gamma', \gamma}(s) J(\gamma')_c \end{aligned}$$

By (??), for each  $\gamma'$ , write

$$(16.2)(b) \quad J(\gamma')_c = \sum_{\delta' \leq \gamma'} (-1)^{\ell(\gamma') - \ell(\delta')} s^{\frac{\ell_0(\gamma') - \ell_0(\delta')}{2}} P_{\delta', \gamma'}(s) I(\delta')_c$$

This formula is computed at (possibly singular, non-integral) infinitesimal character. Plug it in to give

$$(16.2)(c) \quad \begin{aligned} I(\gamma)_c &= I(x, \lambda, (1 - \epsilon)\nu)_c \\ &+ (1 - s) \sum_{\substack{\delta' \leq \gamma' < \gamma \\ \ell(\gamma) - \ell(\gamma') \text{ odd}}} (-1)^{\ell(\gamma') - \ell(\delta')} s^{\frac{\ell_0(\gamma') - \ell_0(\delta')}{2}} P_{\delta', \gamma'}(s) Q_{\gamma', \gamma}(s) I(\delta')_c \end{aligned}$$

or, spelling it out more explicitly:

(16.2)(d)

$$\begin{aligned} I(\gamma)_c &= I(x, \lambda, (1 - \epsilon)\nu)_c \\ &+ (1 - s) \sum_{\substack{\delta' < \gamma \\ \delta' \leq \gamma' < \gamma \\ \ell(\gamma) - \ell(\gamma') \text{ odd}}} s^{\frac{\ell_0(\gamma) - \ell_0(\delta')}{2}} \left[ \sum_{\substack{\gamma' \\ \delta' \leq \gamma' < \gamma \\ \ell(\gamma) - \ell(\gamma') \text{ odd}}} (-1)^{\ell(\gamma') - \ell(\delta')} P_{\delta', \gamma'}(s) Q_{\gamma', \gamma}(s) \right] I(\delta')_c \end{aligned}$$

## 17 Invariant Forms

Suppose  $J$  is a representation of  $Sp(4, \mathbb{R})$  with a central character, and  $\mu$  is a  $K$ -type of  $\pi$ . Identify  $\mu$  with its highest weight  $(r, s)$ .

The element  $\tau$  defining  $Sp(4, \mathbb{R})$  is  $\text{diag}(i, i, -i, -i)$ , which is central in  $K$ , and has square  $-I$ . It acts in  $\mu$  by the scalar  $i^{r+s}$ . Note that if  $-I$  acts in  $J$  by  $\epsilon = \pm 1$  then  $(i^{r+s})^2 = \epsilon$ , i.e.

$$r + s \text{ is } \begin{cases} \text{even} & \epsilon = 1 \\ \text{odd} & \epsilon = -1. \end{cases}$$

Now assume  $J$  is irreducible and  $\mu = (r, s)$  is a lowest  $K$ -type of  $J$ . Write  $\langle, \rangle$  for the *invariant* form on  $J$ , and  $\langle, \rangle_c$  for the c-invariant form as usual.

**Lemma 17.1** *Suppose we have a formula as in (??)*

$$(17.2) \quad J(x, \lambda, \nu)_c = \sum_{x', \lambda'} a(x', \lambda') I_K(x', \lambda')_c.$$

*Suppose  $-I$  acts by  $\epsilon$  in  $J(x, \lambda, \nu)$  and choose  $\zeta^2 = \epsilon$ . Write the (unique) lowest  $K$ -type of  $I_K(x', \lambda')$  as  $(r(x', \lambda'), s(x', \lambda'))$ . Define*

$$\delta(x', \lambda') = \begin{cases} 1 & \zeta i^{r(x', \lambda') + s(x', \lambda')} = 1 \\ s & \zeta i^{r(x', \lambda') + s(x', \lambda')} = -1. \end{cases}$$

*Then an invariant form on  $J$  is given by*

$$(17.3) \quad J(x, \lambda, \nu)_0 = \sum_{x', \lambda'} \delta(x', \lambda') a(x', \lambda') I_K(x', \lambda')_0.$$

*where  $I_K(x', \lambda')_0$  is the unique positive definite invariant form. There is one other invariant form,  $-$  this one.*

*In particular an invariant form on  $J$  is definite if and only if, for all  $x', \lambda'$  appearing in (??),*

$$(17.4)(a) \quad \delta(x', \lambda') a(x', \lambda') \in \mathbb{Z} \text{ for all } (x', \lambda')$$

*or*

$$(17.4)(b) \quad \delta(x', \lambda') a(x', \lambda') \in s\mathbb{Z} \text{ for all } (x', \lambda')$$

## 17.1 Some Invariant Forms on Irreducibles for $Sp(4, \mathbb{R})$

We have some formulas for  $c$ -invariant forms on some irreducible representations: (??)(h,k,n) on  $J(x_9, 3, 1)$ ,  $J(x_7, (\bar{1}, 2), (1, 0))$  and  $J(x_8, (\bar{1}, 2), (1, 0))$ , all at  $\rho$ . Let's convert these to invariant forms.

These representations all have trivial central character. Therefore, we may take  $\zeta = 1$  in Lemma ???. If  $(r, s)$  is a lowest  $K$ -type then the sign in the lemma is  $(-1)^{\frac{r+s}{2}}$ .

First consider  $J(x_8, (\bar{1}, 2), (1, 0))$ .

Equation (??)(n) says

(17.1.5)(a)

$$\boxed{J(x_8, (\bar{1}, 2), (1, 0))_c = I_K(x_8, (\bar{1}, 2))_c - s[I_K(x_1, (2, 1))_c + I_K(x_3, (2, 1))_c]}$$

By Section ?? the lowest  $K$ -types on the right hand side are  $(-1, -3)$ ,  $(1, -3)$  and  $(-3, -3)$ , respectively, which have  $(-1)^{\frac{r+s}{2}} = 1, -1, -1$ , respectively. Therefore

(17.1.5)(b)

$$J(x_8, (\bar{1}, 2), (1, 0))_0 = I_K(x_8, (\bar{1}, 2))_0 - [I_K(x_1, (2, 1))_0 + I_K(x_3, (2, 1))_0].$$

This representation is unitary. This is representation 6 from the output of `block`.

Of course, using (??)(k),  $J(x_7, (\bar{1}, 2), (1, 0))_0$  is essentially the same:

(17.1.5)(c)

$$J(x_7, (\bar{1}, 2), (1, 0))_0 = I_K(x_7, (\bar{1}, 2))_0 - [I_K(x_0, (2, 1))_0 + I_K(x_2, (2, 1))_0].$$

and this is unitary. This is representation 5 from the output of `block`.

Finally (??)(h) says

$$(17.1.5)(d) \quad J(x_9, 3, 1)_c = I_K(x_9, 3) - s[I_K(x_0, (2, 1)) + I_K(x_1, (2, 1))]$$

and the invariant formula is

$$(17.1.5)(e) \quad J(x_9, 3, 1)_0 = I_K(x_9, 3)_0 - [I_K(x_0, (2, 1))_0 + I_K(x_1, (2, 1))_0]$$

which is unitary. This is representation 4 from the output of `block`.

**Remark 17.1.6** *Using the `blocku` command we can confirm the unitarity. These representations are indeed  $A_q(\lambda)$  modules attached to theta stable parabolic subalgebras with Levi factors  $L = U(1) \times SL(2, \mathbb{R})$  and  $L = U(1, 1)$ , respectively.*

## 18 c-Invariant Forms on Irreducible Representations

We already have some formulas for c-invariant forms on some irreducible representations: (??)(h,k,n) on  $J(x_9, 3, 1)$ ,  $J(x_7, (\bar{1}, 2), (1, 0))$  and  $J(x_8, (\bar{1}, 2), (1, 0))$ , all at  $\rho$ .

## 18.1 The c-invariant form on $J(x_9, 1, 3)$

Let's use (??)(p) to get a formula for  $J(x_9, 1, 3)_c$ . Recall  $J(x_9, 1, 3) = J(x_9, (1, 0), \frac{1}{2}(3, 3))$ , irreducible representation 9 from the output of **block** (see Section ??). The formula for  $J(x_9, 1, 3)_c$  is given by (??)(b) with  $a = 2, b = 1$ :

(18.1.1)

$$\begin{aligned} J(x_9, 1, 3)_c = & I(x_9, 1, 3)_c \\ & + I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_2, (2, 1))_c + I(x_3, (2, 1))_c \\ & - I(x_9, 3, 1)_c \\ & - I(x_7, (\bar{1}, 2), (1, 0))_c - I(x_8, (\bar{1}, 2), (1, 0))_c \end{aligned}$$

The terms on the right hand side are given by (??)(p), (??) (4 times), (??)(h) and (??)(d,e), respectively. The last three are also given in (??)(a,d,e). This gives:

(18.1.2)

$$\begin{aligned} J(x_9, 1, 3)_c = & \{I_K(x_9, 1)_c + (1 - s)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c] \\ & + (1 - s)\{2I_K(x_0, (2, 1))_c + 2I_K(x_1, (2, 1))_c\} \\ & + (1 - s)\{I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c\} \\ & + (1 - s)\{I_K(x_9, 3)_c + I_K(x_7, \bar{1}, 2)_c + I_K(x_8, \bar{1}, 2)_c\}\} \\ & + I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c + I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c \\ & - \{I_K(x_9, 3)_c + (1 - s)[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c]\} \\ & - \{I_K(x_7, (\bar{1}, 2))_c + (1 - s)[I_K(x_0, (2, 1))_c + I_K(x_2, (2, 1))_c]\} \\ & - \{I_K(x_8, (\bar{1}, 2))_c + (1 - s)[I_K(x_1, (2, 1))_c + I_K(x_3, (2, 1))_c]\} \end{aligned}$$

Here is a table of the terms:

| $k$ | $\lambda$      | coefficients                       | total   | $(r, s)$   | $\delta$ |
|-----|----------------|------------------------------------|---------|------------|----------|
| 9   | 1              | 1                                  | 1       | $(1, -1)$  | 1        |
| 0   | $(1, 0)$       | $1 - s$                            | $1 - s$ | $(2, 0)$   | $s$      |
| 1   | $(1, 0)$       | $1 - s$                            | $1 - s$ | $(0, -2)$  | $s$      |
| 0   | $(2, 1)$       | $2(1 - s) + 1 - (1 - s) - (1 - s)$ | 1       | $(3, -1)$  | $s$      |
| 1   | $(2, 1)$       | $2(1 - s) + 1 - (1 - s) - (1 - s)$ | 1       | $(1, -3)$  | $s$      |
| 2   | $(2, 1)$       | $(1 - s) + 1 - (1 - s)$            | 1       | $(3, 3)$   | $s$      |
| 3   | $(2, 1)$       | $(1 - s) + 1 - (1 - s)$            | 1       | $(-3, -3)$ | $s$      |
| 9   | 3              | $(1 - s) - 1$                      | $-s$    | $(2, -2)$  | 1        |
| 7   | $(\bar{1}, 2)$ | $(1 - s) - 1$                      | $-s$    | $(3, 1)$   | 1        |
| 8   | $(\bar{1}, 2)$ | $(1 - s) - 1$                      | $-s$    | $(-1, -3)$ | 1        |

And the answer is:

(18.1.3)

$$\begin{aligned}
J(x_9, 1, 3)_c &= I_K(x_9, 1)_c + \\
&\quad + (1 - s) [I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c] \\
&\quad + [I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c + I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c] \\
&\quad - s [I_K(x_9, 3)_c + I_K(x_7, (\bar{1}, 2))_c + I_K(x_8, (\bar{1}, 2))_c]
\end{aligned}$$

Note that if  $s = 1$  this gives (??)(b) again, as it must.

Using Section ??, the invariant form is given as follows.

(18.1.4)

$$\begin{aligned}
J(x_9, 1, 3)_0 &= I_K(x_9, 1)_0 + \\
&\quad + (s - 1) [I_K(x_0, (1, 0))_0 + I_K(x_1, (1, 0))_0] \\
&\quad + s [I_K(x_0, (2, 1))_0 + I_K(x_1, (2, 1))_0 + I_K(x_2, (2, 1))_0 + I_K(x_3, (2, 1))_0] \\
&\quad - s [I_K(x_9, 3)_0 + I_K(x_7, (\bar{1}, 2))_0 + I_K(x_8, (\bar{1}, 2))_0]
\end{aligned}$$

This is not unitary. Note that we can tell this from the previous expression because of the hyperbolic term  $(1 - s)$ .

## 18.2 The c-invariant form on $J(x_{7,8}, (\bar{2}, 1), (2, 0))$

The character formula for  $J(x_7, (\bar{2}, 1), (2, 0))$  is (??)(e), and this holds as a formula for c-invariant forms:



$$\begin{aligned}
(18.2.5) \quad J(x_7, (\bar{2}, 1), (2, 0))_c &= I(x_7, (\bar{2}, 1), (2, 0))_c \\
&+ I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_2, (2, 1))_c \\
&- I(x_9, (3, 1))_c - I(x_7, (\bar{1}, 2), (1, 0))_c
\end{aligned}$$

The non-discrete series terms on the right hand side are given by (??)(d), (??)(h) and (??)(c). The result is:

(18.2.6)

$$\begin{aligned}
J(x_7, (\bar{2}, 1), (2, 0))_c &= -s[I(x_9, 3)_c + I(x_7, (\bar{1}, 2))_c] \\
&+ [I(x_0, (1, 0))_c + I(x_2, (1, 0))_c] \\
&+ [I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_2, (2, 1))_c]
\end{aligned}$$

The lowest  $K$ -types of these terms are:  $(2, -2), (3, 1), (2, 0), (2, 2), (3, -1), (1, -3), (3, 3)$ . We multiply by  $1, 1, s, 1, s, s, s$  respectively, to give the invariant form:

(18.2.7)

$$\begin{aligned}
J(x_7, (\bar{2}, 1), (2, 0))_0 &= -s[I(x_9, 3)_0 + I(x_7, (\bar{1}, 2))_0] \\
&+ [sI(x_0, (1, 0))_0 + I(x_2, (1, 0))_0] \\
&+ s[I(x_0, (2, 1))_0 + I(x_1, (2, 1))_0 + I(x_2, (2, 1))_0]
\end{aligned}$$

This representation is *not* unitary: the signs differ on the two lowest  $K$ -types  $(2, 2)$  and  $(2, 0)$ .

Similarly:

(18.2.8)

$$\begin{aligned}
J(x_8, (\bar{2}, 1), (2, 0))_c &= -s[I(x_9, 3)_c + I(x_8, (\bar{1}, 2))_c] \\
&+ [I(x_1, (1, 0))_c + I(x_3, (1, 0))_c] \\
&+ [I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_3, (2, 1))_c]
\end{aligned}$$

## 19 The Trivial Representation

Let's prove the trivial representation  $J(x_{10}, (\bar{2}, \bar{1}), (2, 1))$  is unitary.

The deformation will go as follows. Let  $\nu = (1 + \epsilon)(2, 1)$  for  $\epsilon > 0$  small. Deforming  $\nu$  to 0 we have the following reducibility points, up to small deformations.

$$(19.1)(a) \quad \nu = (2, 1), \left(\frac{4}{3}, \frac{2}{3}\right), \left(1, \frac{1}{2}\right), \left(\frac{2}{3}, \frac{1}{3}\right)$$

At the first point the integral root system is  $C_2$ ; at all the others it is of type  $A_1$ .

Here is the reducibility at  $\nu = (\frac{2}{3}, \frac{1}{3})$ , obtained using `nblock`.

The input is  $\lambda = \rho = (2, 1)$ , so  $\lambda - \rho = (0, 0)$ .

```
real: nblock
choose Cartan class (one of 0,1,2,3): 3
Choosing the unique KGB element for the Cartan class:
10:  3  [r,r]   10  10      *   *   (0,0)#3 1^2x1^e
rho = [1,1]/1
Give lambda-rho: 0 0
denominator for nu: 3
numerator for nu: 1 1
x=10, lambda=[1,1]/1, gamma=[1,1]/3.
Name an output file (return for stdout, ? to abandon):
Subsystem on dual side is of type A1, with roots 6.
Given parameters define element 1 of the following block:
0(0,2):  0  [i2]  0   (1,2)  *(x= 4, nu= [2,-1]/6,,lam=rho+ [-2,0])  2,1,2
1(1,0):  1  [r2]  2   (0,*)  *(x=10, nu=  [1,1]/3,,lam=rho+[-2,-2])  e
2(1,1):  1  [r2]  1   (0,*)  *(x=10, nu=  [1,1]/3,,lam=rho+[-2,-1])  e
KL polynomials (-1)^(l(1)-l(x))*P_{x,1}:
0: -1
1: 1
```

This says:

$$(19.1)(b) \quad I(x_{10}, (\bar{2}, \bar{1}), (\frac{2}{3}, \frac{1}{3})) = J(x_{10}, (\bar{2}, \bar{1}), (\frac{2}{3}, \frac{1}{3})) + J(x_9, 1, \frac{1}{3}).$$

**Dangerous Bend:** Here come some orientation numbers.

Applying (??) as usual, for the first time we have some nontrivial orientation numbers.

By Table ??4,  $(\bar{a} = 0, \bar{b} = 1, y = \frac{1}{3})$ ,  $\ell_0(x_{10}, (\bar{2}, \bar{1}), (\frac{2}{3}, \frac{1}{3})) = 3$ . On the other hand by Table (??).2, with  $(c = 1, x = \frac{1}{3})$ ,  $\ell_0(x_9, 1, \frac{1}{3}) = 1$ . The orientation number contribution to (??) is  $s^{\frac{1}{2}(3-1)} = s$ , so:

$$(19.1)(c) \quad I(x_{10}, (\bar{2}, \bar{1}), (\frac{2}{3}, \frac{1}{3}))_c = I_K(x_{10}, (\bar{2}, \bar{1}))_c + (1-s)sI_K(x_9, 1)_c \\ = I_K(x_{10}, (\bar{2}, \bar{1}))_c + (s-1)I_K(x_9, 1)_c.$$

The next reducibility point is  $(1, \frac{1}{2})$ . We're interested in  $I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))$ . This is of type  $A_{sn}^I$  in the notation of [?, Section 6]. Here is **nblock**:

```
real: nblock
choose Cartan class (one of 0,1,2,3): 3
Choosing the unique KGB element for the Cartan class:
10: 3  [r,r]  10  10      *   *   (0,0)#3 1^2x1^e
rho = [1,1]/1
Give lambda-rho: 0 0
denominator for nu: 2
numerator for nu: 1 1
x=10, lambda=[1,1]/1, gamma=[1,1]/2.
Name an output file (return for stdout, ? to abandon):
Subsystem on dual side is of type A1, with roots 7.
Given parameters define element 2 of the following block:
0(0,1): 0  [i1]  1  (2,*)  *(x= 5, nu= [-1,1]/2,,lam=rho+ [0,-1])  1,2,1
1(1,1): 0  [i1]  0  (2,*)  *(x= 6, nu= [-1,1]/2,,lam=rho+ [0,-1])  1,2,1
2(2,0): 1  [r1]  2  (0,1)  *(x=10, nu=  [1,1]/2,,lam=rho+[-2,-2])  e
KL polynomials (-1)^(l(2)-l(x))*P_{x,2}:
0: -1
1: -1
2: 1
```

This says

$$(19.1)(d) \quad I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2})) = J(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2})) + J(x_5, (1, \bar{0}), (0, \frac{1}{2})) + J(x_6, (1, \bar{0}), (0, \frac{1}{2}))$$

or alternatively

$$(19.1)(e) \quad I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2})) = J(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2})) + J(x_7, (\bar{0}, 1), (\frac{1}{2}, 0)) + J(x_8, (\bar{0}, 1), (\frac{1}{2}, 0))$$

Now apply (??). There is an orientation number here:  $\ell_0(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2})) = 2$  (Table ??1) and  $\ell_0(x_{7,8}, (\bar{0}, 1), (\frac{1}{2}, 0)) = 0$  (Table ??1). So:

$$(19.1)(f) \quad I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))_c = I(x_{10}, (\bar{2}, \bar{1}), (\frac{2}{3}, \frac{1}{3}))_c \\ + (1-s)s[I_K(x_7, (\bar{0}, 1))_c + I_K(x_8, (\bar{0}, 1))_c]$$

Plugging in (c) gives

$$(19.1)(g) \quad I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))_c = I_K(x_{10}, (\bar{2}, \bar{1}))_c + (s-1)I_K(x_9, 1)_c \\ + (s-1)[I_K(x_7, (\bar{0}, 1))_c + I_K(x_8, (\bar{0}, 1))_c]$$

However there is one more step:  $I(x_7, (\bar{0}, 1))$  and  $I(x_8, (\bar{0}, 1))$  are not final.

Use (??) to give

$$(19.1)(h) \quad I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))_c = I_K(x_{10}, (\bar{2}, \bar{1}))_c + (s-1)I_K(x_9, 1)_c \\ + (s-1)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c + I_K(x_2, (1, 0))_c + I_K(x_3, (1, 0))_c]$$

Let's move on to  $(\frac{4}{3}, \frac{2}{3})$ , i.e.  $I(x_{10}, (\bar{2}, \bar{1}), (\frac{4}{3}, \frac{2}{3}))$ . This standard module is irreducible. The integral real root is  $(1, 1)$ . By the table in Section ?? this fails the parity condition, so doesn't give reducibility. This is also what nblock says:

```
real: nblock
choose Cartan class (one of 0,1,2,3): 3
Choosing the unique KGB element for the Cartan class:
10: 3 [r,r] 10 10 * * (0,0)#3 1^2x1^e
rho = [1,1]/1
Give lambda-rho: 0 0
denominator for nu: 3
numerator for nu: 2 2
x=10, lambda=[1,1]/1, gamma=[2,2]/3.
Name an output file (return for stdout, ? to abandon):
Subsystem on dual side is of type A1, with roots 6.
Given parameters define element 0 of the following block:
0(0,0): 0 [rn] 0 (*,*) *(x=10, nu= [2,2]/3, lam=rho+[-2,-2]) e
KL polynomials (-1)^(l(0)-l(x))*P_{x,0}:
0: 1
```

Note that in [?, Section 6], case  $A_{sn}^I$ , this example doesn't appear (meaning it is irreducible); in that notation we'd be considering  $J(x, \bar{2}e, (\frac{4}{3}, \frac{2}{3}))$ , but what appears in the table is  $J(x, \bar{2}o, (\frac{4}{3}, \frac{2}{3}))$ .

Finally we come to  $\nu = (2, 1)$ . By (??)(j) with  $a = 2, b = 1$ :

$$\begin{aligned}
 (19.1)(i) \quad I(x_{10}, (\bar{2}, \bar{1}), (2, 1))^3 = & \\
 & J(x_{10}, (\bar{2}, \bar{1}), (2, 1))^3 \\
 & + J(x_0, (2, 1))^0 + J(x_1, (2, 1))^0 \\
 & + 2 \times J(x_9, 3, 1)^1 \\
 & + J(x_7, (\bar{1}, 2), (1, 0))^1 + J(x_8, (\bar{1}, 2), (1, 0))^1 \\
 & + J(x_7, (\bar{2}, 1), (2, 0))^2 + J(x_8, (\bar{2}, 1), (2, 0))^2 \\
 & + J(x_9, 1, 3)^2
 \end{aligned}$$

Remember  $J(x_9, c, x) = J(x_9, (c, 0), \frac{1}{2}(x, x))$ , with infinitesimal character  $\frac{1}{2}(x + c, x - c)$ .

We're at integral infinitesimal character, so there are no orientation numbers, so (??) gives

$$\begin{aligned}
 (19.1)(j) \quad I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c = & I(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))_c \\
 & + (1 - s) [I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c] \\
 & + (1 - s) [J(x_7, (\bar{2}, 1), (2, 0))_c + J(x_8, (\bar{2}, 1), (2, 0))_c + J(x_9, 1, 3)_c]
 \end{aligned}$$

and plugging in (h) for  $J(x_{10}, (\bar{2}, \bar{1}), (1, \frac{1}{2}))_c$  gives

$$\begin{aligned}
 (19.1)(k) \quad I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c = & I_K(x_{10}, (\bar{2}, \bar{1}))_c \\
 & + (s - 1)I_K(x_9, 1)_c \\
 & + (s - 1) [I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c + I_K(x_2, (1, 0))_c + I_K(x_3, (1, 0))_c] \\
 & + (1 - s) [I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c] \\
 & + (1 - s) [J(x_7, (\bar{2}, 1), (2, 0))_c + J(x_8, (\bar{2}, 1), (2, 0))_c + J(x_9, 1, 3)_c]
 \end{aligned}$$

We still have to deal with the terms  $J(x_7, (\bar{2}, 1), (2, 0))$   $J(x_8, (\bar{2}, 1), (2, 0))$  and  $J(x_9, 1, 3)$ .

These are available from (??), (??) and (??). So:

$$\begin{aligned}
(19.1)(1) \\
I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c &= I_K(x_{10}, (\bar{2}, \bar{1}))_c \\
&+ (s-1)I_K(x_9, 1)_c \\
&+ (s-1)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c + I_K(x_2, (1, 0))_c + I_K(x_3, (1, 0))_c] \\
&+ (1-s)[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c] \\
\\
&+ (1-s)\{ \\
&- s[I(x_9, 3)_c + I(x_7, (\bar{1}, 2))_c] \\
&+ I(x_0, (1, 0))_c + I(x_2, (1, 0))_c \\
&+ I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_2, (2, 1))_c \\
\\
&- s[I(x_9, 3)_c + I(x_8, (\bar{1}, 2))_c] \\
&+ I(x_1, (1, 0))_c + I(x_3, (1, 0))_c \\
&+ I(x_0, (2, 1))_c + I(x_1, (2, 1))_c + I(x_3, (2, 1))_c \\
\\
&+ I_K(x_9, 1)_c + (1-s)[I_K(x_0, (1, 0))_c + I_K(x_1, (1, 0))_c] \\
&+ I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c + I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c \\
&- s[I_K(x_9, 3)_c + I_K(x_7, (\bar{1}, 2))_c + I_K(x_8, \bar{1}, 2)_c] \quad \}
\end{aligned}$$

Here is a table of the terms:

| $k$ | $\lambda$            | coefficients                    | total    |
|-----|----------------------|---------------------------------|----------|
| 10  | $(\bar{2}, \bar{1})$ | 1                               | 1        |
| 9   | 1                    | $(s-1) + (1-s)$                 | 0        |
| 9   | 3                    | $(1-s) + (1-s) + (1-s)$         | $3(1-s)$ |
| 8   | $(\bar{1}, 2)$       | $(1-s) + (1-s)$                 | $2(1-s)$ |
| 7   | $(\bar{1}, 2)$       | $(1-s) + (1-s)$                 | $2(1-s)$ |
| 3   | $(2, 1)$             | $(1-s) + (1-s)$                 | $2(1-s)$ |
| 2   | $(2, 1)$             | $(1-s) + (1-s)$                 | $2(1-s)$ |
| 1   | $(2, 1)$             | $(1-s) + (1-s) + (1-s) + (1-s)$ | $4(1-s)$ |
| 0   | $(2, 1)$             | $(1-s) + (1-s) + (1-s) + (1-s)$ | $4(1-s)$ |
| 3   | $(1, 0)$             | $(s-1) + (1-s)$                 | 0        |
| 2   | $(1, 0)$             | $(s-1) + (1-s)$                 | 0        |
| 1   | $(1, 0)$             | $(s-1) + (1-s) + (1-s)^2$       | $2(1-s)$ |
| 0   | $(1, 0)$             | $(s-1) + (1-s) + (1-s)^2$       | $2(1-s)$ |

The character formula for the trivial representation is (??)(j) with  $a = 2, b = 1$ , and this holds without change for c-invariant forms (see Section ??):

$$\begin{aligned}
(19.1)(m) \quad & J(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c = I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c \\
& - \{I(x_9, 1, 3)_c + I(x_7, (\bar{2}, 1), (2, 0))_c + I(x_8, (\bar{2}, 1), (2, 0))_c\} \\
& + \{I(x_7, (\bar{1}, 2), (1, 0))_c + I(x_8, (\bar{1}, 2), (1, 0))_c + I(x_9, 3, 1)_c\} \\
& - \{I(x_3, (2, 1))_c + I(x_2, (2, 1))_c + I(x_1, (2, 1))_c + I(x_0, (2, 1))_c\}
\end{aligned}$$

We know the terms on the right hand side:

- (1)  $I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c$ : (??)
- (2)  $I(x_9, 1, 3)_c$ : (??)
- (3)  $I(x_9, 3, 1)_c$ : (??)(h)
- (4)  $I(x_k, (\bar{1}, 2), (1, 0))_c$  ( $k = 7, 8$ ): (??)
- (5)  $I(x_k, (\bar{2}, 1), (2, 0))_c$  ( $k = 7, 8$ ): (??)(d) and (e)

Of course the last four terms require no further comment.

So let's tabulate everything. The next table has a column for each standard module on the right hand side of (??), each row is a limit  $K$ -representation  $I_K(x, \lambda)$ , and the columns give the multiplicities in the expression of the c-invariant form on the standard module.

|                                     | $I(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c$ | $I(x_9, 1, 3)_c$ | $I(x_7, (\bar{2}, 1), (2, 0))_c$ | $I(x_8, (\bar{2}, 1), (2, 0))_c$ |
|-------------------------------------|-------------------------------------------|------------------|----------------------------------|----------------------------------|
| $I_K(x_{10}, (\bar{2}, \bar{1}))_c$ | 1                                         |                  |                                  |                                  |
| $I_K(x_9, 1)_c$                     | 0                                         | -1               |                                  |                                  |
| $I_K(x_9, 3)_c$                     | $3(1-s)$                                  | $-(1-s)$         | $-(1-s)$                         | $-(1-s)$                         |
| $I_K(x_8, (\bar{1}, 2))_c$          | $2(1-s)$                                  | $-(1-s)$         |                                  | $-(1-s)$                         |
| $I_K(x_7, (\bar{1}, 2))_c$          | $2(1-s)$                                  | $-(1-s)$         | $-(1-s)$                         |                                  |
| $I_K(x_3, (2, 1))_c$                | $2(1-s)$                                  | $-(1-s)$         |                                  | $-(1-s)$                         |
| $I_K(x_2, (2, 1))_c$                | $2(1-s)$                                  | $-(1-s)$         | $-(1-s)$                         |                                  |
| $I_K(x_1, (2, 1))_c$                | $4(1-s)$                                  | $-2(1-s)$        | $-(1-s)$                         | $-2(1-s)$                        |
| $I_K(x_0, (2, 1))_c$                | $4(1-s)$                                  | $-2(1-s)$        | $-2(1-s)$                        | $-(1-s)$                         |
| $I_K(x_3, (1, 0))_c$                |                                           |                  |                                  | -1                               |
| $I_K(x_2, (1, 0))_c$                |                                           |                  | -1                               |                                  |
| $I_K(x_1, (1, 0))_c$                | $2(1-s)$                                  | $-(1-s)$         |                                  | -1                               |
| $I_K(x_0, (1, 0))_c$                | $2(1-s)$                                  | $-(1-s)$         | -1                               |                                  |

|                                     | $I(x_7, (\bar{1}, 2), (1, 0))_c$ | $I(x_8, (\bar{1}, 2), (1, 0))_c$ | $I(x_9, 3, 1)_c$ | DS | Sum  |
|-------------------------------------|----------------------------------|----------------------------------|------------------|----|------|
| $I_K(x_{10}, (\bar{2}, \bar{1}))_c$ |                                  |                                  |                  |    | 1    |
| $I_K(x_9, 1)_c$                     |                                  |                                  |                  |    | -1   |
| $I_K(x_9, 3)_c$                     |                                  |                                  | 1                |    | 1    |
| $I_K(x_8, (\bar{1}, 2))_c$          |                                  | 1                                |                  |    | 1    |
| $I_K(x_7, (\bar{1}, 2))_c$          | 1                                |                                  |                  |    | 1    |
| $I_K(x_3, (2, 1))_c$                |                                  | $(1-s)$                          |                  | -1 | $-s$ |
| $I_K(x_2, (2, 1))_c$                | $(1-s)$                          |                                  |                  | -1 | $-s$ |
| $I_K(x_1, (2, 1))_c$                |                                  | $(1-s)$                          | $(1-s)$          | -1 | $-s$ |
| $I_K(x_0, (2, 1))_c$                | $(1-s)$                          |                                  | $(1-s)$          | -1 | $-s$ |
| $I_K(x_3, (1, 0))_c$                |                                  |                                  |                  |    | -1   |
| $I_K(x_2, (1, 0))_c$                |                                  |                                  |                  |    | -1   |
| $I_K(x_1, (1, 0))_c$                |                                  |                                  |                  |    | $-s$ |
| $I_K(x_0, (1, 0))_c$                |                                  |                                  |                  |    | $-s$ |

So:

(19.1)(n)

$$\begin{aligned}
J(x_{10}, (\bar{2}, \bar{1}), (2, 1))_c &= I_K(x_{10}, (\bar{2}, \bar{1}))_c \\
&+ [I_K(x_9, 3)_c - I_K(x_9, 1)_c] \\
&+ [I_K(x_7, (\bar{1}, 2))_c + I_K(x_8, (\bar{1}, 2))_c] \\
&- s[I_K(x_0, (2, 1))_c + I_K(x_1, (2, 1))_c + I_K(x_2, (2, 1))_c + I_K(x_3, (2, 1))_c] \\
&- [sI_K(x_0, (1, 0))_c + sI_K(x_1, (1, 0))_c + I_K(x_2, (1, 0))_c + I_K(x_3, (1, 0))_c]
\end{aligned}$$



To get the invariant form, we give the lowest  $K$ -types and the corresponding correction factors:

|                                   |            |     |
|-----------------------------------|------------|-----|
| $I_K(x_{10}, (\bar{2}, \bar{1}))$ | $(0, 0)$   | 1   |
| $I_K(x_9, 1)$                     | $(1, -1)$  | 1   |
| $I_K(x_9, 3)$                     | $(2, -2)$  | 1   |
| $I_K(x_8, (\bar{1}, 2))$          | $(-1, -3)$ | 1   |
| $I_K(x_7, (\bar{1}, 2))$          | $(3, 1)$   | 1   |
| $I_K(x_3, (2, 1))$                | $(-3, -3)$ | $s$ |
| $I_K(x_2, (2, 1))$                | $(3, 3)$   | $s$ |
| $I_K(x_1, (2, 1))$                | $(1, -3)$  | $s$ |
| $I_K(x_0, (2, 1))$                | $(3, -1)$  | $s$ |
| $I_K(x_3, (1, 0))$                | $(-2, -2)$ | 1   |
| $I_K(x_2, (1, 0))$                | $(2, 2)$   | 1   |
| $I_K(x_1, (1, 0))$                | $(0, -2)$  | $s$ |
| $I_K(x_0, (1, 0))$                | $(2, 0)$   | $s$ |

Taking this into account we compute the invariant form on the trivial representation  $J(x_{10}, (\bar{2}, \bar{1}), (2, 1))$ :

$$\begin{aligned}
(19.1)(o) \\
J(x_{10}, (\bar{2}, \bar{1}), (2, 1))_0 &= I_K(x_{10}, (\bar{2}, \bar{1})) \\
&+ [I_K(x_9, 3) - I_K(x_9, 1)] \\
&+ [I_K(x_7, (\bar{1}, 2)) + I_K(x_8, (\bar{1}, 2))] \\
&- [I_K(x_0, (2, 1)) + I_K(x_1, (2, 1)) + I_K(x_2, (2, 1)) + I_K(x_3, (2, 1))] \\
&- [I_K(x_0, (1, 0)) + I_K(x_1, (1, 0)) + I_K(x_2, (1, 0)) + I_K(x_3, (1, 0))].
\end{aligned}$$

Therefore the trivial representation of  $Sp(4, \mathbb{R})$  is unitary. If anyone knows an easier proof please let me know.

## References

- [1] Jeffrey Adams Some notes on parametrizing representations. [www.liegroups.org/papers/basepoint.pdf](http://www.liegroups.org/papers/basepoint.pdf).
- [2] Annegret Paul Assigning Representation Parameters to Atlas Block Output. [www.liegroups.org/papers/Dictionary08.pdf](http://www.liegroups.org/papers/Dictionary08.pdf).

- [3] David A. Vogan, Jr. Branching example:  $Sp(4, \mathbb{R})$ . [www-math.mit.edu/dav/sp4ex.pdf](http://www-math.mit.edu/dav/sp4ex.pdf).
- [4] David A. Vogan, Jr. Forms example:  $Sp(4, \mathbb{R})$ . [www.liegroups.org/papers/sp4forms.pdf](http://www.liegroups.org/papers/sp4forms.pdf).