

**Sample Problems for the First In-Class Exam**  
**Math 246, Spring 2018, Professor David Levermore**

- (1) (a) Give the integral being evaluated by the following Matlab command.

```
int('x/(1+x^4)', 'x', 0, inf)
```

- (b) Sketch the graph that would be produced by the following Matlab command.

```
ezplot('2/t', [1, 6])
```

- (c) Sketch the graph that would be produced by the following Matlab commands.

```
[X, Y] = meshgrid(-5:0.1:5, -5:0.1:5)  
contour(X, Y, X.^2 + Y.^2, [1, 9, 25])  
axis square
```

- (2) Find the explicit solution for each of the following initial-value problems and identify its interval of definition.

(a)  $\frac{dz}{dt} = \frac{\cos(t) - z}{1 + t}, \quad z(0) = 2.$

(b)  $\frac{du}{dz} = e^u + 1, \quad u(0) = 0.$

(c)  $\frac{dv}{dt} = -3t^2e^{-v}, \quad v(2) = 0.$

- (3) Give the interval of definition for the solution of the initial-value problem

$$\frac{dx}{dt} + \frac{1}{t^2 - 4}x = \frac{1}{\sin(t)}, \quad x(1) = 0.$$

(You do not have to solve this equation to answer this question!)

- (4) Consider the following Matlab commands.

```
>> [T, Y] = meshgrid(-5.0:1.0:5.0, -5.0:1.0:5.0);  
>> S = T.^2 - Y.^3;  
>> L = sqrt(1 + S.^2);  
>> quiver(T, Y, 1./L, S./L, 0.5)  
>> axis tight, xlabel 't', ylabel 'y'
```

- (a) What is the differential equation being studied?

- (b) What kind of graph will these Matlab commands produce?

(5) Consider the differential equation

$$\frac{dy}{dt} = \frac{y^2(y+2)(y-4)}{y-2}.$$

- (a) Sketch its phase-line portrait over the interval  $[-6, 6]$ . Identify points where it is undefined. Identify its stationary points and classify each as being either stable, unstable, or semistable.
  - (b) For each stationary point identify the set of initial values  $y(0)$  such that the solution  $y(t)$  converges to that stationary point as  $t \rightarrow \infty$ .
  - (c) For each stationary point identify the set of initial values  $y(0)$  such that the solution  $y(t)$  converges to that stationary point as  $t \rightarrow -\infty$ .
  - (d) Identify all initial values  $y(0)$  such that the interval of definition of the solution  $y(t)$  is  $(-\infty, \infty)$ .
  - (e) Sketch a graph of  $y$  versus  $t$  showing several solution curves. The graph should show all of the stationary solutions as well as solution curves above and below each of them. Every value of  $y$  for which the equation is defined should lie on at least one sketched solution curve.
- (6) In the absence of predators the population of mosquitoes in a certain area would increase at a rate proportional to its current population such that it would triple every five weeks. There are 85,000 mosquitoes in the area when a flock of birds arrives that eats 25,000 mosquitoes per week. Write down an initial-value problem that governs  $M(t)$ , the population of mosquitoes in the area after the flock of birds arrives. (You do not have to solve the initial-value problem!)
- (7) A tank initially contains 100 liters of pure water. Beginning at time  $t = 0$  brine (salt water) with a salt concentration of 2 grams per liter (gr/lit) flows into the tank at a constant rate of 3 liters per minute (lit/min) and the well-stirred mixture flows out of the tank at the same rate. Let  $S(t)$  denote the mass (gr) of salt in the tank at time  $t \geq 0$ .
- (a) Write down an initial-value problem that governs  $S(t)$ .
  - (b) Is  $S(t)$  an increasing or decreasing function of  $t$ ? (Give your reasoning.)
  - (c) What is the behavior of  $S(t)$  as  $t \rightarrow \infty$ ? (Give your reasoning.)
  - (d) Derive an explicit formula for  $S(t)$ .
  - (e) How does the answer to part (a) change if the well-stirred mixture flows out of the tank at a constant rate of 2 liters per minute?
- (8) A 2 kilogram (kg) mass initially at rest is dropped in a medium that offers a resistance of  $v^2/40$  newtons ( $= \text{kg m/sec}^2$ ) where  $v$  is the downward velocity (m/sec) of the mass. The gravitational acceleration is  $9.8 \text{ m/sec}^2$ .
- (a) What is the terminal velocity of the mass?
  - (b) Write down an initial-value problem that governs  $v$  as a function of time. (You do not have to solve it!)

(9) Give an implicit general solution to each of the following differential equations.

(a)  $\left(\frac{y}{x} + 3x\right) dx + (\log(x) - y) dy = 0.$

(b)  $(x^2 + y^3 + 2x) dx + 3y^2 dy = 0.$

(10) Suppose we are using the Runge-midpoint method to numerically approximate the solution of an initial-value problem over the time interval  $[0, 5]$ . By what factor would we expect the error to decrease when we increase the number of time steps taken from 500 to 2000?

(11) Consider the following Matlab function m-file.

```
function [t,y] = solveit(tI, yI, tF, n)

t = zeros(n + 1, 1); y = zeros(n + 1, 1);
t(1) = tI; y(1) = yI; h = (tF - tI)/n;
for i = 1:n
z = t(i)^4 + y(i)^2;
t(i + 1) = t(i) + h;
y(i + 1) = y(i) + (h/2)*(z + t(i + 1)^4 + (y(i) + h*z)^2);
end
```

Suppose the input values are  $tI = 1$ ,  $yI = 1$ ,  $tF = 5$ , and  $n = 20$ .

- What is the initial-value problem being approximated numerically?
- What is the numerical method being used?
- What is the step size?
- What are the output values of  $t(2)$  and  $y(2)$ ?

(12) Suppose we have used a numerical method to approximate the solution of an initial-value problem over the time interval  $[1, 6]$  with 1000 uniform time steps. How many uniform time steps do we need to reduce the global error of our approximation by roughly a factor of  $\frac{1}{81}$  if the method we had used was each of the following?

- Explicit Euler method
- Runge-trapezoidal method
- Runge-midpoint method
- Runge-Kutta method