

Final Exam Sample Problems, Math 246, Spring 2018

- (1) Consider the differential equation $\frac{dy}{dt} = (9 - y^2)y^2$.
- Find all of its stationary points and classify their stability.
 - Sketch its phase-line portrait in the interval $-5 \leq y \leq 5$.
 - If $y_1(0) = -1$, how does the solution $y_1(t)$ behave as $t \rightarrow \infty$?
 - If $y_2(0) = 4$, how does the solution $y_2(t)$ behave as $t \rightarrow \infty$?
 - Evaluate

$$\lim_{t \rightarrow \infty} (y_2(t) - y_1(t)).$$

- (2) Solve each of the following initial-value problems and give the interval of definition of each solution.

(a) $\frac{dy}{dt} + \frac{2ty}{1+t^2} = t^2, \quad y(0) = 1.$

(b) $\frac{dy}{dx} + \frac{e^x y + 2x}{2y + e^x} = 0, \quad y(0) = 0.$

- (3) Determine constants a and b such that the following differential equation is exact. Then find a general solution in implicit form.

$$(ye^x + y^3) dx + (ae^x + bxy^2) dy = 0.$$

- (4) Consider the following Matlab function m-file.

```
function [t,y] = solveit(ti, yi, tf, n)
t = zeros(n + 1, 1); y = zeros(n + 1, 1);
t(1) = ti; y(1) = yi; h = (tf - ti)/n;
for i = 1:n
t(i + 1) = t(i) + h; y(i + 1) = y(i) + h*((t(i))^4 + (y(i))^2);
end
```

Suppose that the input values are $ti = 1$, $yi = 1$, $tf = 5$, and $n = 40$.

- What initial-value problem is being approximated numerically?
 - What numerical method is being used?
 - What is the step size?
 - What are the output values of $t(2)$, $y(2)$, $t(3)$, and $y(3)$?
- (5) Consider the following Matlab commands.

```
[t,y] = ode45(@(t,y) y.*(y-1).*(2-y), [0,3], -0.5:0.5:2.5); plot(t,y)
```

The following questions need not be answered in Matlab format!

- What is the differential equation being solved numerically?
- Give the initial condition for each solution being approximated?
- Over what time interval are the solutions being approximated?
- Sketch each of these solutions over this time interval on a single graph. Label the initial value of each solution clearly.
- What is the limiting behavior of each solution as $t \rightarrow \infty$?

(6) Let $y(t)$ be the solution of the initial-value problem

$$y' = 4t(y + y^2), \quad y(0) = 1.$$

- (a) Use two steps of the explicit Euler method to approximate $y(1)$.
 (b) Use one step of the Runge-trapezoidal method to approximate $y(1)$.
 (c) Use one step of the Runge-midpoint method to approximate $y(1)$.

(7) Give an explicit real-valued general solution of the following equations.

(a) $y'' - 2y' + 5y = te^t + \cos(2t)$

(b) $\ddot{u} - 3\dot{u} - 10u = te^{-2t}$

(c) $v'' + 9v = \cos(3t)$

(d) $w'''' + 13w'' + 36w = 9\sin(t)$

(8) Given that $y_1(t) = t$ and $y_2(t) = t^{-2}$ are solutions of the associated homogeneous equation, find a general solution of

$$t^2 y'' + 2t y' - 2y = \frac{3}{t^2} + 5t, \quad \text{for } t > 0.$$

(9) Solve the following initial-value problems.

(a) $w'' + 4w' + 20w = 5e^{2t}, \quad w(0) = 3, \quad w'(0) = -7.$

(b) $y'' - 4y' + 4y = \frac{e^{2t}}{3+t}, \quad y(0) = 0, \quad y'(0) = 5.$

Evaluate any definite integrals that arise.

(10) Give an explicit real-valued general solution of the equation

$$\ddot{h} + 2\dot{h} + 5h = 0.$$

Sketch a typical solution for $t \geq 0$. If this equation governs a spring-mass system, is the system undamped, under damped, critically damped, or over damped? (Give your reasoning!)

(11) When a mass of 2 kilograms is hung vertically from a spring, it stretches the spring 0.5 m. (Gravitational acceleration is 9.8 m/sec².) At $t = 0$ the mass is set in motion from 0.3 meters below its rest (equilibrium) position with a upward velocity of 2 m/sec. It is acted upon by an external force of $2\cos(5t)$. Neglect damping and assume that the spring force is proportional to its displacement. Formulate an initial-value problem that governs the motion of the mass for $t > 0$. (Do not solve this initial-value problem; just write it down!)

(12) Find the Laplace transform $Y(s)$ of the solution $y(t)$ to the initial-value problem

$$y'' + 4y' + 8y = f(t), \quad y(0) = 2, \quad y'(0) = 4.$$

where

$$f(t) = \begin{cases} 4 & \text{for } 0 \leq t < 2, \\ t^2 & \text{for } 2 \leq t. \end{cases}$$

You may refer to the table of Laplace transforms on the last page. (Do not take the inverse Laplace transform to find $y(t)$; just solve for $Y(s)$!)

(13) Find the function $y(t)$ whose Laplace transform $Y(s)$ is given by

$$(a) \quad Y(s) = \frac{e^{-3s}4}{s^2 - 6s + 5}, \quad (b) \quad Y(s) = \frac{e^{-2s}s}{s^2 + 4s + 8}.$$

You may refer to the table of Laplace transforms on the last page.

(14) Two interconnected tanks, each with a capacity of 60 liters, contain brine (salt water). At $t = 0$ the first tank contains 22 liters and the second contains 17 liters. Brine with a salt concentration of 6 grams per liter flows into the first tank at 7 liters per hour. Well-stirred brine flows from the first tank into the second at 8 liters per hour, from the second into the first at 5 liters per hour, from the first into a drain at 2 liter per hour, and from the second into a drain at 4 liters per hour. At $t = 0$ there are 31 grams of salt in the first tank and 43 grams in the second.

- Determine the volume of brine in each tank as a function of time.
- Give an initial-value problem that governs the amount of salt in each tank as a function of time. (Do not solve the IVP.)
- Give the interval of definition for the solution of this initial-value problem.

(15) Consider the real vector-valued functions $\mathbf{x}_1(t) = \begin{pmatrix} 1 \\ t \end{pmatrix}$, $\mathbf{x}_2(t) = \begin{pmatrix} t^3 \\ 3 + t^4 \end{pmatrix}$.

- Compute the Wronskian $\text{Wr}[\mathbf{x}_1, \mathbf{x}_2](t)$.
- Find $\mathbf{A}(t)$ such that $\mathbf{x}_1, \mathbf{x}_2$ is a fundamental set of solutions to the linear system $\mathbf{x}' = \mathbf{A}(t)\mathbf{x}$.
- Give a general solution to the system you found in part (b).

(16) Give a real, vector-valued general solution of the linear planar system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ for

$$(a) \quad \mathbf{A} = \begin{pmatrix} 6 & 4 \\ 4 & 0 \end{pmatrix}, \quad (b) \quad \mathbf{A} = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}.$$

(17) Sketch the phase-plane portrait of the linear planar system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ for

$$(a) \quad \mathbf{A} = \begin{pmatrix} 6 & 4 \\ 4 & 0 \end{pmatrix}, \quad (b) \quad \mathbf{A} = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}.$$

(18) What answer will be produced by the following Matlab command?

$$\gg A = [1 \ 4; 3 \ 2]; [\text{vect}, \text{val}] = \text{eig}(\text{sym}(A))$$

You do not have to give the answer in Matlab format.

(19) A real 2×2 matrix \mathbf{B} has the eigenpairs

$$\left(2, \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right) \quad \text{and} \quad \left(-1, \begin{pmatrix} -1 \\ 2 \end{pmatrix} \right).$$

- Give a general solution to the linear planar system $\mathbf{x}' = \mathbf{B}\mathbf{x}$.
- Give an invertible matrix \mathbf{V} and a diagonal matrix \mathbf{D} that diagonalize \mathbf{B} .
- Compute $e^{t\mathbf{B}}$.
- Sketch a phase-plane portrait for this system and identify its type. Classify the stability of the origin. Carefully mark all sketched orbits with arrows!

(20) Solve the initial-value problem $\mathbf{x}' = \mathbf{A}\mathbf{x}$, $\mathbf{x}(0) = \mathbf{x}^I$ for the following \mathbf{A} and \mathbf{x}^I .

(a) $\mathbf{A} = \begin{pmatrix} 3 & 10 \\ -5 & -7 \end{pmatrix}$, $\mathbf{x}^I = \begin{pmatrix} -3 \\ 2 \end{pmatrix}$.

(b) $\mathbf{A} = \begin{pmatrix} 8 & -5 \\ 5 & -2 \end{pmatrix}$, $\mathbf{x}^I = \begin{pmatrix} 3 \\ -1 \end{pmatrix}$.

(21) Consider the system

$$\dot{x} = 2xy, \quad \dot{y} = 9 - 9x - y^2.$$

- Find all of its stationary points.
- Find all of its semistationary orbits.
- Find a nonconstant function $H(x, y)$ such that every orbit of the system satisfies $H(x, y) = c$ for some constant c .
- Classify the type and stability of each stationary point.
- Sketch the stationary points plus the level set $H(x, y) = c$ for each value of c that corresponds to a stationary point that is a saddle. Carefully mark all sketched orbits with arrows!

(22) Consider the system

$$u' = -5v, \quad v' = u - 4v - u^2.$$

- Find all of its stationary points.
- Compute the Jacobian matrix at each stationary point.
- Classify the type and stability of each stationary point.
- Sketch a phase-plane portrait of the system that shows its behavior near each stationary point. Carefully mark all sketched orbits with arrows!

(23) Consider the system

$$\dot{p} = p(3 - 3p + 2q), \quad \dot{q} = q(6 - p - q).$$

- Find all of its stationary points.
- Compute the Jacobian matrix at each stationary point.
- Classify the type and stability of each stationary point.
- Sketch a phase-plane portrait of the system that shows its behavior near each stationary point. Carefully mark all sketched orbits with arrows!
- Add the orbits of all semistationary solutions to the phase-plane portrait sketched for part (d). Carefully mark these sketched orbits with arrows!
- Why do solutions that start in the first quadrant stay in the first quadrant?

Table of Laplace Transforms

$h(t) = \mathcal{L}^{-1}[H](t)$	$H(s) = \mathcal{L}[h](s)$
$t^n e^{at}$ for $n \geq 0$	$\frac{n!}{(s-a)^{n+1}}$ for $s > a$
$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}$ for $s > a$
$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}$ for $s > a$
$e^{at} \cosh(bt)$	$\frac{s-a}{(s-a)^2 - b^2}$ for $s > a + b $
$e^{at} \sinh(bt)$	$\frac{b}{(s-a)^2 - b^2}$ for $s > a + b $
$t^n j(t)$ for $n \geq 0$	$(-1)^n J^{(n)}(s)$ where $J(s) = \mathcal{L}[j](s)$
$j'(t)$	$sJ(s) - j(0)$ where $J(s) = \mathcal{L}[j](s)$
$e^{at} j(t)$	$J(s-a)$ where $J(s) = \mathcal{L}[j](s)$
$u(t-c)j(t-c)$ for $c \geq 0$	$e^{-cs}J(s)$ where $J(s) = \mathcal{L}[j](s)$
$\delta(t-c)j(t)$ for $c \geq 0$	$e^{-cs}j(c)$

Here a , b , and c are real numbers; n is an integer; $j(t)$ is any function that is nice enough; $u(t)$ is the unit step (Heaviside) function; $\delta(t)$ is the unit impulse (Dirac delta).