

Sample Problems for the First In-Class Exam
Math 246, Fall 2018, Professor David Levermore

- (1) (a) Sketch the graph that would be produced by the following Matlab command.

```
fplot(@(t) 2/t, [1,6])
```

- (b) Sketch the graph that would be produced by the following Matlab commands.

```
[X, Y] = meshgrid(-5:0.1:5, -5:0.1:5)  
contour(X, Y, X.^2 + Y.^2, [1, 9, 25])  
axis square
```

- (2) Find the explicit solution for each of the following initial-value problems and identify its interval of definition.

(a) $\frac{dz}{dt} = \frac{\cos(t) - z}{1 + t}, \quad z(0) = 2.$

(b) $\frac{du}{dz} = e^u + 1, \quad u(0) = 0.$

(c) $\frac{dv}{dt} = -3t^2 e^{-v}, \quad v(2) = 0.$

- (3) Give the interval of definition for the solution of the initial-value problem

$$\frac{dx}{dt} + \frac{1}{t^2 - 4} x = \frac{1}{\sin(t)}, \quad x(1) = 0.$$

(You do not need to solve this equation to answer this question, but your reasoning must be given!)

- (4) Consider the following Matlab commands.

```
>> [T, Y] = meshgrid(-5.0:1.0:5.0, -5.0:1.0:5.0);  
>> S = T.^2 - Y.^3;  
>> L = sqrt(1 + S.^2);  
>> quiver(T, Y, 1./L, S./L, 0.5)  
>> axis tight, xlabel 't', ylabel 'y'
```

- (a) What is the differential equation being studied?
(b) What kind of graph will these Matlab commands produce?

(5) Consider the differential equation

$$\frac{dy}{dt} = \frac{y^2(y+2)(y-4)}{y-2}.$$

- (a) Sketch its phase-line portrait over the interval $[-6, 6]$. Identify points where it is undefined. Identify its stationary points and classify each as being either stable, unstable, or semistable.
 - (b) For each stationary point identify the set of initial values $y(0)$ such that the solution $y(t)$ converges to that stationary point as $t \rightarrow \infty$.
 - (c) For each stationary point identify the set of initial values $y(0)$ such that the solution $y(t)$ converges to that stationary point as $t \rightarrow -\infty$.
 - (d) Identify all initial values $y(0)$ such that the interval of definition of the solution $y(t)$ is $(-\infty, \infty)$.
 - (e) Sketch a graph of y versus t showing several solution curves. The graph should show all of the stationary solutions as well as solution curves above and below each of them. Every value of y for which the equation is defined should lie on at least one sketched solution curve.
- (6) In the absence of predators the population of mosquitoes in a certain area would increase at a rate proportional to its current population such that it would triple every five weeks. There are 85,000 mosquitoes in the area when a flock of birds arrives that eats 25,000 mosquitoes per week. Write down an initial-value problem that governs $M(t)$, the population of mosquitoes in the area after the flock of birds arrives. (You do not have to solve the initial-value problem!)
- (7) A tank initially contains 100 liters of pure water. Beginning at time $t = 0$ brine (salt water) with a salt concentration of 2 grams per liter (gr/lit) flows into the tank at a constant rate of 3 liters per minute (lit/min) and the well-stirred mixture flows out of the tank at the same rate. Let $S(t)$ denote the mass (gr) of salt in the tank at time $t \geq 0$.
- (a) Write down an initial-value problem that governs $S(t)$.
 - (b) Is $S(t)$ an increasing or decreasing function of t ? (Give your reasoning.)
 - (c) What is the behavior of $S(t)$ as $t \rightarrow \infty$? (Give your reasoning.)
 - (d) Derive an explicit formula for $S(t)$.
 - (e) How does the answer to part (a) change if the well-stirred mixture flows out of the tank at a constant rate of 2 liters per minute?
- (8) A 2 kilogram (kg) mass initially at rest is dropped in a medium that offers a resistance of $v^2/40$ newtons ($= \text{kg m/sec}^2$) where v is the downward velocity (m/sec) of the mass. The gravitational acceleration is 9.8 m/sec^2 .
- (a) What is the terminal velocity of the mass?
 - (b) Write down an initial-value problem that governs v as a function of time. (You do not have to solve it!)

(9) Give an implicit general solution to each of the following differential equations.

(a) $\left(\frac{y}{x} + 3x\right) dx + (\log(x) - y) dy = 0.$

(b) $(x^2 + y^3 + 2x) dx + 3y^2 dy = 0.$

(10) Suppose we are using the Runge-midpoint method to numerically approximate the solution of an initial-value problem over the time interval $[0, 5]$. By what factor would we expect the error to decrease when we increase the number of time steps taken from 500 to 2000?

(11) Consider the following Matlab function m-file.

```
function [t,y] = solveit(tI, yI, tF, n)

t = zeros(n + 1, 1); y = zeros(n + 1, 1);
t(1) = tI; y(1) = yI; h = (tF - tI)/n;
for i = 1:n
z = t(i)^4 + y(i)^2;
t(i + 1) = t(i) + h;
y(i + 1) = y(i) + (h/2)*(z + t(i + 1)^4 + (y(i) + h*z)^2);
end
```

Suppose the input values are $tI = 1$, $yI = 1$, $tF = 5$, and $n = 20$.

- What is the initial-value problem being approximated numerically?
- What is the numerical method being used?
- What is the step size?
- What are the output values of $t(2)$ and $y(2)$?

(12) Suppose we have used a numerical method to approximate the solution of an initial-value problem over the time interval $[1, 6]$ with 1000 uniform time steps. How many uniform time steps do we need to reduce the global error of our approximation by roughly a factor of $\frac{1}{81}$ if the method we had used was each of the following?

- Explicit Euler method
- Runge-trapezoidal method
- Runge-midpoint method
- Runge-Kutta method