

Final Exam Sample Problems, Math 246, Fall 2019

- (1) Consider the differential equation $\frac{dy}{dt} = (9 - y^2)y^2$.
- Find all of its stationary points and classify their stability.
 - Sketch its phase-line portrait in the interval $-5 \leq y \leq 5$.
 - If $y_1(0) = -1$, how does the solution $y_1(t)$ behave as $t \rightarrow \infty$?
 - If $y_2(0) = 4$, how does the solution $y_2(t)$ behave as $t \rightarrow \infty$?
 - Evaluate

$$\lim_{t \rightarrow \infty} (y_2(t) - y_1(t)).$$

- (2) Solve each of the following initial-value problems and give the interval of definition of each solution.

(a) $x' = \frac{t}{(t^2 + 1)x}, \quad x(0) = -3.$

(b) $\frac{dy}{dt} + \frac{2ty}{1 + t^2} = t^2, \quad y(0) = 1.$

(c) $\frac{dy}{dx} + \frac{e^x y + 2x}{2y + e^x} = 0, \quad y(0) = 0.$

- (3) Determine constants a and b such that the following differential equation is exact. Then find a general solution in implicit form.

$$(ye^x + y^3) dx + (ae^x + bxy^2) dy = 0.$$

- (4) Consider the following Matlab function m-file.

```
function [t,y] = solveit(ti, yi, tf, n)
t = zeros(n + 1, 1); y = zeros(n + 1, 1);
t(1) = ti; y(1) = yi; h = (tf - ti)/n;
for i = 1:n
t(i + 1) = t(i) + h; y(i + 1) = y(i) + h*((t(i))^4 + (y(i))^2);
end
```

Suppose that the input values are $t_i = 1$, $y_i = 1$, $t_f = 5$, and $n = 40$.

- What initial-value problem is being approximated numerically?
 - What numerical method is being used?
 - What is the step size?
 - What are the output values of $t(2)$, $y(2)$, $t(3)$, and $y(3)$?
- (5) Let $y(t)$ be the solution of the initial-value problem

$$y' = 4t(y + y^2), \quad y(0) = 1.$$

- Use two steps of the explicit Euler method to approximate $y(1)$.
- Use one step of the Runge-trapezoidal method to approximate $y(1)$.
- Use one step of the Runge-midpoint method to approximate $y(1)$.

- (6) Consider the following Matlab commands.

```
[t,y] = ode45(@(t,y) y.*(y-1).*(2-y), [0,3], -0.5:0.5:2.5); plot(t,y)
```

The following questions need not be answered in Matlab format!

- What is the differential equation being solved numerically?
- Give the initial condition for each solution being approximated?
- Over what time interval are the solutions being approximated?
- Sketch each of these solutions over this time interval on a single graph. Label the initial value of each solution clearly.
- What is the limiting behavior of each solution as $t \rightarrow \infty$?

- (7) Suppose we are using the Runge-midpoint method to numerically approximate the solution of an initial-value problem over the time interval $[1, 9]$. By what factor would we expect the error to decrease when we increase the number of time steps taken from 400 to 2000?

- (8) A NASA engineer has used the Runge-Kutta method to approximate the solution of an initial-value problem over the time interval $[2, 10]$ with 800 uniform time steps.

- How many uniform time steps are needed to reduce the global error by a factor of $\frac{1}{256}$?
- What is the order of a numerical method that reduces the global error by a factor of $\frac{1}{256}$ when the step size is halved?

- (9) Give an explicit real-valued general solution of the following equations.

- $y'' - 2y' + 5y = t e^t + \cos(2t)$
- $\ddot{u} - 3\dot{u} - 10u = t e^{-2t}$
- $v'' + 9v = \cos(3t)$
- $w'''' + 13w'' + 36w = 9 \sin(t)$

- (10) Solve the following initial-value problems.

- $w'' + 4w' + 20w = 5e^{2t}$, $w(0) = 3$, $w'(0) = -7$.
- $y'' - 4y' + 4y = \frac{e^{2t}}{3+t}$, $y(0) = 0$, $y'(0) = 5$.

Evaluate any definite integrals that arise.

- (11) Given that $y_1(t) = t$ and $y_2(t) = t^{-2}$ solve the associated homogeneous equation, find a general solution of

$$t^2 y'' + 2t y' - 2y = \frac{3}{t^2} + 5t, \quad \text{for } t > 0.$$

- (12) Given that t^2 and $t^2 \log(t)$ solve the associated homogeneous differential equation, consider the initial-value problem

$$t^2 x'' - 3t x' + 4x = t^2 \log(t), \quad x(1) = 0, \quad x'(1) = 0.$$

- Give the interval of definition of its solution.
- Compute $\text{Wr}[t^2, t^2 \log(t)]$.
- Find $x(t)$. Evaluate any definite integrals that arise.

- (13) Give an explicit real-valued general solution of the equation

$$\ddot{h} + 2\dot{h} + 5h = 0.$$

Sketch a typical solution for $t \geq 0$. If this equation governs a spring-mass system, is the system undamped, under damped, critically damped, or over damped? (Give your reasoning!)

- (14) When a mass of 2 kilograms is hung vertically from a spring, it stretches the spring 0.5 m. (Gravitational acceleration is 9.8 m/sec^2 .) At $t = 0$ the mass is set in motion from 0.3 meters below its rest (equilibrium) position with a upward velocity of 2 m/sec. It is acted upon by an external force of $2 \cos(5t)$. Neglect damping and assume that the spring force is proportional to its displacement. Formulate an initial-value problem that governs the motion of the mass for $t > 0$. (Do not solve this initial-value problem; just write it down!)

- (15) Find the Laplace transform $Y(s)$ of the solution $y(t)$ to the initial-value problem

$$y'' + 4y' + 8y = f(t), \quad y(0) = 2, \quad y'(0) = 4.$$

where

$$f(t) = \begin{cases} 4 & \text{for } 0 \leq t < 2, \\ t^2 & \text{for } 2 \leq t. \end{cases}$$

You may refer to the table of Laplace transforms on the last page. (Do not take the inverse Laplace transform to find $y(t)$; just solve for $Y(s)$!)

- (16) Let $x(t)$ be the solution of the initial-value problem

$$x'' + 10x' + 29x = f(t), \quad x(0) = 3, \quad x'(0) = -7,$$

where the forcing $f(t)$ is given by

$$f(t) = \begin{cases} t^2 & \text{for } 0 \leq t < 1, \\ e^{1-t} & \text{for } 1 \leq t < \infty. \end{cases}$$

(a) Find the Laplace transform $F(s)$ of the forcing $f(t)$.

(b) Find the Laplace transform $X(s)$ of the solution $x(t)$.

(DO NOT take the inverse Laplace transform to find $x(t)$; just solve for $X(s)$!)

You may refer to the table of Laplace transforms on the last page.

- (17) Find the function $y(t)$ whose Laplace transform $Y(s)$ is given by

$$(a) \quad Y(s) = \frac{e^{-3s}4}{s^2 - 6s + 5}, \quad (b) \quad Y(s) = \frac{e^{-2s}s}{s^2 + 4s + 8}.$$

You may refer to the table of Laplace transforms on the last page.

- (18) Consider the real vector-valued functions $\mathbf{x}_1(t) = \begin{pmatrix} 1 \\ t \end{pmatrix}$, $\mathbf{x}_2(t) = \begin{pmatrix} t^3 \\ 3 + t^4 \end{pmatrix}$.

(a) Compute the Wronskian $\text{Wr}[\mathbf{x}_1, \mathbf{x}_2](t)$.

(b) Find $\mathbf{A}(t)$ such that $\mathbf{x}_1, \mathbf{x}_2$ is a fundamental set of solutions to the linear system $\mathbf{x}' = \mathbf{A}(t)\mathbf{x}$.

(c) Give a general solution to the system you found in part (b).

- (19) Two interconnected tanks, each with a capacity of 60 liters, contain brine (salt water). At $t = 0$ the first tank contains 22 liters and the second contains 17 liters. Brine with a salt concentration of 6 grams per liter flows into the first tank at 7 liters per hour. Well-stirred brine flows from the first tank into the second at 8 liters per hour, from the second into the first at 5 liters per hour, from the first into a drain at 2 liter per hour, and from the second into a drain at 4 liters per hour. At $t = 0$ there are 31 grams of salt in the first tank and 43 grams in the second.
- Determine the volume of brine in each tank as a function of time.
 - Give an initial-value problem that governs the amount of salt in each tank as a function of time. (Do not solve the IVP.)
 - Give the interval of definition for the solution of this initial-value problem.

- (20) Give a real, vector-valued general solution of the linear planar system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ for

$$(a) \quad \mathbf{A} = \begin{pmatrix} 6 & 4 \\ 4 & 0 \end{pmatrix}, \quad (b) \quad \mathbf{A} = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}.$$

- (21) Sketch the phase-plane portrait of the linear planar system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ for

$$(a) \quad \mathbf{A} = \begin{pmatrix} 6 & 4 \\ 4 & 0 \end{pmatrix}, \quad (b) \quad \mathbf{A} = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}.$$

- (22) What answer will be produced by the following Matlab command?

```
>> A = [1 4; 3 2]; [vect, val] = eig(sym(A))
```

You do not have to give the answer in Matlab format.

- (23) A real 2×2 matrix \mathbf{B} has the eigenpairs

$$\left(2, \begin{pmatrix} 3 \\ 1 \end{pmatrix}\right) \quad \text{and} \quad \left(-1, \begin{pmatrix} -1 \\ 2 \end{pmatrix}\right).$$

- Give a general solution to the linear planar system $\mathbf{x}' = \mathbf{B}\mathbf{x}$.
- Give an invertible matrix \mathbf{V} and a diagonal matrix \mathbf{D} that diagonalize \mathbf{B} .
- Compute $e^{t\mathbf{B}}$.
- Find \mathbf{B} .
- Sketch a phase-plane portrait for this system and identify its type. Classify the stability of the origin. Carefully mark all sketched orbits with arrows!

- (24) Solve the initial-value problem $\mathbf{x}' = \mathbf{A}\mathbf{x}$, $\mathbf{x}(0) = \mathbf{x}^I$ for the following \mathbf{A} and \mathbf{x}^I .

$$(a) \quad \mathbf{A} = \begin{pmatrix} 3 & 10 \\ -5 & -7 \end{pmatrix}, \quad \mathbf{x}^I = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$

$$(b) \quad \mathbf{A} = \begin{pmatrix} 8 & -5 \\ 5 & -2 \end{pmatrix}, \quad \mathbf{x}^I = \begin{pmatrix} 3 \\ -1 \end{pmatrix}.$$

$$(c) \quad \mathbf{A} = \begin{pmatrix} -2 & 1 \\ -1 & -4 \end{pmatrix}, \quad \mathbf{x}^I = \begin{pmatrix} 3 \\ 1 \end{pmatrix}.$$

(25) Consider the system

$$\dot{x} = 2xy, \quad \dot{y} = 9 - 9x - y^2.$$

- Find all of its stationary points.
- Find all of its semistationary orbits.
- Find a nonconstant function $H(x, y)$ such that every orbit of the system satisfies $H(x, y) = c$ for some constant c .
- Classify the type and stability of each stationary point.
- Sketch the stationary points plus the level set $H(x, y) = c$ for each value of c that corresponds to a stationary point that is a saddle. Carefully mark all sketched orbits with arrows!

(26) Consider the system

$$\dot{p} = -9p + 3q, \quad \dot{q} = 4p - 8q + 10p^2.$$

- This system has two stationary points. Find them.
- Find the Jacobian matrix at each stationary point.
- Classify the type and stability of each stationary point.
- Sketch a phase-plane portrait of the system that shows its behavior near each stationary point. Carefully mark all sketched orbits with arrows!

(27) Consider the system

$$u' = -5v, \quad v' = u - 4v - u^2.$$

- Find all of its stationary points.
- Compute the Jacobian matrix at each stationary point.
- Classify the type and stability of each stationary point.
- Sketch a phase-plane portrait of the system that shows its behavior near each stationary point. Carefully mark all sketched orbits with arrows!

(28) Consider the system

$$\dot{p} = p(3 - 3p + 2q), \quad \dot{q} = q(6 - p - q).$$

- Find all of its stationary points.
- Compute the Jacobian matrix at each stationary point.
- Classify the type and stability of each stationary point.
- Sketch a phase-plane portrait of the system that shows its behavior near each stationary point. Carefully mark all sketched orbits with arrows!
- Add the orbits of all semistationary solutions to the phase-plane portrait sketched for part (d). Carefully mark these sketched orbits with arrows!
- Why do solutions that start in the first quadrant stay in the first quadrant?

Table of Laplace Transforms

	$h(t) = \mathcal{L}^{-1}[H](t)$	$H(s) = \mathcal{L}[h](s)$
1.	$t^n e^{at}$ for $n \geq 0$	$\frac{n!}{(s-a)^{n+1}}$ for $s > a$
2.	$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}$ for $s > a$
3.	$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}$ for $s > a$
4.	$e^{at} \cosh(bt)$	$\frac{s-a}{(s-a)^2 - b^2}$ for $s > a + b $
5.	$e^{at} \sinh(bt)$	$\frac{b}{(s-a)^2 - b^2}$ for $s > a + b $
6.	$t^n j(t)$ for $n \geq 0$	$(-1)^n J^{(n)}(s)$ where $J(s) = \mathcal{L}[j](s)$
7.	$j'(t)$	$sJ(s) - j(0)$ where $J(s) = \mathcal{L}[j](s)$
8.	$e^{at} j(t)$	$J(s-a)$ where $J(s) = \mathcal{L}[j](s)$
9.	$u(t-c)j(t-c)$ for $c \geq 0$	$e^{-cs}J(s)$ where $J(s) = \mathcal{L}[j](s)$
10.	$\delta(t-c)j(t)$ for $c \geq 0$	$e^{-cs}j(c)$

Here a , b , and c are real numbers; n is an integer; $j(t)$ is any function that is nice enough; $u(t)$ is the unit step (Heaviside) function; $\delta(t)$ is the unit impulse (Dirac delta).